



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/25, C07K 14/545, 16/24, C12N 1/21, 5/10, 15/62, C07K 17/00, A61K 39/44, 39/395, G01N 33/68, C12Q 1/68, C07K 17/14	A2	(11) International Publication Number: WO 00/17363 (43) International Publication Date: 30 March 2000 (30.03.00)
(21) International Application Number: PCT/US99/20868 (22) International Filing Date: 17 September 1999 (17.09.99) (30) Priority Data: 09/156,966 18 September 1998 (18.09.98) US (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US). (72) Inventor: TIMANS, Jacqueline, C.; 1538 Canna Court, Mountain View, CA 94043 (US). (74) Agents: GOULD, James, M. et al.; Schering-Plough Corporation, Patent Dept., K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: INTERLEUKIN-1 LIKE MOLECULE: INTERLEUKIN-1-ZETA; NUCLEIC ACIDS, POLYPEPTIDES, ANTIBODIES, THEIR USES (57) Abstract Nucleic acids encoding mammalian, e.g., primate, IL-1 ζ , purified IL-1 ζ polypeptides and fragments thereof. Binding proteins, e.g., antibodies, both polyclonal and monoclonal, are also provided. Methods of using the compositions for both diagnostic and therapeutic utilities are provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERLEUKIN-1 LIKE MOLECULE: INTERLEUKIN-1-ZETA; NUCLEIC ACIDS, POLYPEPTIDES, ANTIBODIES, THEIR USES

This filing is a PCT Patent Application claiming priority to U.S. Patent Application USSN 09/156,966, filed September 18, 1998.

5

FIELD OF THE INVENTION

The present invention relates to compositions and methods for affecting mammalian physiology, including morphogenesis or immune system function. In particular, it provides nucleic
10 acids, proteins, and antibodies which regulate development and/or the immune system. Diagnostic and therapeutic uses of these materials are also disclosed.

BACKGROUND OF THE INVENTION

15 Recombinant DNA technology refers generally to techniques of integrating genetic information from a donor source into vectors for subsequent processing, such as through introduction into a host, whereby the transferred genetic information is copied and/or expressed in the new environment.
20 Commonly, the genetic information exists in the form of complementary DNA (cDNA) derived from messenger RNA (mRNA) coding for a desired protein product. The carrier is frequently a plasmid having the capacity to incorporate cDNA for later replication in a host and, in some cases, actually
25 to control expression of the cDNA and thereby direct synthesis of the encoded product in the host.

For some time, it has been known that the mammalian immune response is based on a series of complex cellular interactions, called the "immune network". Recent research
30 has provided new insights into the inner workings of this network. While it remains clear that much of the immune response does, in fact, revolve around the network-like interactions of lymphocytes, macrophages, granulocytes, and other cells, immunologists now generally hold the opinion that
35 soluble proteins, known as lymphokines, cytokines, or monokines, play critical roles in controlling these cellular interactions. Thus, there is considerable interest in the

isolation, characterization, and mechanisms of action of cell modulatory factors, an understanding of which will lead to significant advancements in the diagnosis and therapy of numerous medical abnormalities, e.g., immune system disorders.

5 Lymphokines apparently mediate cellular activities in a variety of ways. They have been shown to support the proliferation, growth, and/or differentiation of pluripotent hematopoietic stem cells into vast numbers of progenitors comprising diverse cellular lineages which make up a complex
10 immune system. Proper and balanced interactions between the cellular components are necessary for a healthy immune response. The different cellular lineages often respond in a different manner when lymphokines are administered in conjunction with other agents.

15 Cell lineages especially important to the immune response include two classes of lymphocytes: B-cells, which can produce and secrete immunoglobulins (proteins with the capability of recognizing and binding to foreign matter to effect its removal), and T-cells of various subsets that
20 secrete lymphokines and induce or suppress the B-cells and various other cells (including other T-cells) making up the immune network. These lymphocytes interact with many other cell types.

Another important cell lineage is the mast cell (which
25 has not been positively identified in all mammalian species), which is a granule-containing connective tissue cell located proximal to capillaries throughout the body. These cells are found in especially high concentrations in the lungs, skin, and gastrointestinal and genitourinary tracts. Mast cells
30 play a central role in allergy-related disorders, particularly anaphylaxis as follows: when selected antigens crosslink one class of immunoglobulins bound to receptors on the mast cell surface, the mast cell degranulates and releases mediators, e.g., histamine, serotonin, heparin, and prostaglandins, which
35 cause allergic reactions, e.g., anaphylaxis.

Research to better understand and treat various immune disorders has been hampered by the general inability to

maintain cells of the immune system in vitro. Immunologists have discovered that culturing many of these cells can be accomplished through the use of T-cell and other cell supernatants, which contain various growth factors, including
5 many of the lymphokines.

The interleukin-1 family of proteins includes the IL-1 α , the IL-1 β , the IL-1RA, and recently the IL-1 γ (also designated Interferon-Gamma Inducing Factor, IGIF). This related family of genes have been implicated in a broad range of biological
10 functions. See Dinarello (1994) FASEB J. 8:1314-1325; Dinarello (1991) Blood 77:1627-1652; and Okamura, et al. (1995) Nature 378:88-91.

In addition, various growth and regulatory factors exist which modulate morphogenetic development. This includes,
15 e.g., the Toll ligands, which signal through binding to receptors which share structural, and mechanistic, features characteristic of the IL-1 receptors. See, e.g., Lemaitre; et al. (1996) Cell 86:973-983; and Belvin and Anderson (1996) Ann. Rev. Cell & Develop. Biol. 12:393-416.

20 From the foregoing, it is evident that the discovery and development of new soluble proteins, including ones similar to lymphokines, should contribute to new therapies for a wide range of degenerative or abnormal conditions which directly or indirectly involve development, differentiation, or function,
25 e.g., of the immune system and/or hematopoietic cells. In particular, the discovery and understanding of novel lymphokine-like molecules which enhance or potentiate the beneficial activities of other lymphokines would be highly advantageous. The present invention provides new interleukin-
30 1 like compositions and related compounds, and methods for their use.

SUMMARY OF THE INVENTION

The present invention is based on the discovery, purification, and characterization of the biological activities of a novel mammalian, e.g., primate, interleukin-1 like molecule, designated interleukin-1 ζ (IL-1 ζ). IL-1 ζ exhibits both structural and sequence similarity, e.g., by homology comparison, to known members of the IL-1 family of molecules.

In a first aspect, the invention provides an isolated or recombinant polypeptide that: specifically binds polyclonal antibodies generated against at least a 12 consecutive amino acid segment of SEQ ID NO: 2 or 4; and comprises at least one sequence selected from: GENSGVK; EDWEKD; CCLEDPA; FVHTSR; KKFSIHD; VLVLDS; NLIAVP; FFALAS; SSASAEK; SLILLGV; FCLYCDK; PSLQLK; KLMKLAAQ; FIFYRAQ; SRNMLES; WFICTS; EPVGVT; or FSFQPVC (see SEQ ID NO: 2); or FVHTSP; SPILLGV; or SWNMLES (see SEQ ID NO: 4). Certain embodiments include those: wherein the polypeptide comprises a plurality of the sequence; or which specifically bind to polyclonal antibodies generated against an immunogen selected from SEQ ID NO: 2 or 4. Other embodiments include those where the 12 consecutive amino acid segment is selected from: GVKMGSEDWEKD; AGSPLEPGPSLP; SRKVKSLNPKKF; HDQDHKVLVLDS; NLIAVPDKNYIR; FALASSLSSASA; GQSHPSLQLKKE; MKLAAQKESARR; FYRAQVGSRNML; TSCNCNEPVGVT; FENRKHIEFSFQ; or PVCKAEMSPSEV (see SEQ ID NO: 2); or AVSPLEPGPSLP; SPKVKNLNPCKKF; or FYRAQVGSWNML (see SEQ ID NO: 4). Certain preferred embodiments include those wherein the polypeptide: comprises a mature protein; lacks a post-translational modification; is from a primate, including a human; is a natural allelic variant of IL-1 ζ ; has a length at least about 30 amino acids; exhibits at least two non-overlapping epitopes that are specific for a primate IL-1 ζ ; exhibits a sequence identity over a length of at least about 20 amino acids to SEQ ID NO: 2 or 4; is not glycosylated; has a molecular weight of at least 10 kD with natural glycosylation; is a synthetic polypeptide; is attached to a solid substrate; is conjugated to another chemical moiety; is

a 5-fold or less substitution from natural sequence; or is a deletion or insertion variant from a natural sequence.

Other embodiments include a soluble polypeptide comprising: the sterile polypeptide; the sterile polypeptide
5 and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

Fusion protein embodiments include those having a polypeptide sequence described, further comprising: a mature
10 polypeptide as described; a detection or purification tag, including a FLAG, His6, or Ig sequence; or sequence of another cytokine or chemokine.

Kit embodiments include those comprising such a polypeptide and: a compartment comprising said polypeptide;
15 and/or instructions for use or disposal of reagents in said kit.

Antibody or binding compound embodiments encompass a binding compound comprising an antigen binding site from an antibody, which specifically binds to a mature polypeptide, as
20 described, wherein: the mature polypeptide is a primate IL-1 ζ ; the binding compound is an Fv, Fab, or Fab2 fragment; the binding compound is conjugated to another chemical moiety; or the antibody: is raised against a 12 consecutive amino acid segment of SEQ ID NO: 2 or 4; is raised against a mature IL-
25 1 ζ ; is raised to a purified primate IL-1 ζ ; is immunoselected; is a polyclonal antibody; binds to a denatured IL-1 ζ ; exhibits a Kd to antigen of at least 30 μ M; is attached to a solid substrate, including a bead or plastic membrane; is in a sterile composition; or is detectably labeled, including a
30 radioactive or fluorescent label. An alternative binding compound embraces one comprising an antigen binding portion from an antibody, which specifically binds to a primate protein, as described, wherein: the protein is a human protein; the binding compound is an Fv, Fab, or Fab2 fragment;
35 the binding compound is conjugated to another chemical moiety; or the antibody: is raised against a polypeptide sequence of a mature polypeptide comprising at least 12 consecutive amino

acids of SEQ ID NO: 2 or 4; is raised against a mature primate IL-1 ζ ; is raised to a purified primate IL-1 ζ ; is immunoselected; is a polyclonal antibody; binds to a denatured primate IL-1 ζ ; exhibits a Kd to antigen of at least 30 μ M; is
5 attached to a solid substrate, including a bead or plastic membrane; is in a sterile composition; or is detectably labeled, including a radioactive or fluorescent label. Kits are provided comprising the binding compound, as described, and: a compartment comprising said binding compound; and/or
10 instructions for use or disposal of reagents in the kit. Methods are also provided, e.g., of: making an antibody, comprising immunizing an immune system with an immunogenic amount of: a primate IL-1 ζ polypeptide; or a peptide sequence comprising at least 12 consecutive amino acids of SEQ ID NO: 2
15 or 4; thereby causing said antibody to be produced; or producing an antigen:antibody complex, comprising contacting a primate IL-1 ζ polypeptide with an antibody, as described, thereby allowing said complex to form.

The invention further embraces a composition comprising:
20 the sterile binding compound described, or the binding compound and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

25 Nucleic acid embodiments include an isolated or recombinant nucleic acid encoding the described polypeptide, wherein: the polypeptide is a primate IL-1 ζ ; or the nucleic acid: encodes an antigenic peptide sequence of SEQ ID NO: 2 or 4; encodes a plurality of antigenic peptide sequences of SEQ
30 ID NO: 2 or 4; exhibits at least about 80% identity to a natural cDNA encoding said segment; is an expression vector; further comprises an origin of replication; is from a natural source; comprises a detectable label; comprises synthetic nucleotide sequence; is less than 6 kb, preferably less than 3
35 kb; is from a rodent; comprises a natural full length coding sequence; is a hybridization probe for a gene encoding said IL-1 ζ ; or xiii) is a PCR primer, PCR product, or mutagenesis

primer; or encodes an IL-1 ζ polypeptide. The invention also provides a cell transformed with the described nucleic acid, e.g., where the cell is: a prokaryotic cell; a eukaryotic cell; a bacterial cell; a yeast cell; an insect cell; a
5 mammalian cell; a mouse cell; a primate cell; or a human cell.

Another embodiment is a kit comprising the described nucleic acid and: a compartment comprising said nucleic acid and: a compartment further comprising a primate IL-1 ζ polypeptide; and/or instructions for use or disposal of
10 reagents in said kit.

Other nucleic acid embodiments include an isolated or recombinant nucleic acid that hybridizes under wash conditions of 30° C and less than 2M salt to SEQ ID NO: 1; or where the wash condition is at: 45° C and/or 500 mM salt; 55° C and/or
15 150 mM salt; or encodes at least 12 or 17 contiguous amino acids of SEQ ID NO: 2 or 4.

The invention also provides methods of modulating a cell involved in an inflammatory response comprising contacting said cell with an agonist or antagonist of a primate IL-1 ζ
20 polypeptide. Preferably, the contacting is in combination with an agonist or antagonist of IL-1 α , IL-1RA, IL-1 β , IL-1 γ , IL-1 δ , IL-1 ϵ , IL-2, and/or IL-12; the contacting is with an antagonist, including binding composition comprising an antibody binding site which specifically binds an IL-1 ζ ; or
25 the modulating is regulation of IFN- γ production.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

OUTLINE

- I. General
- 5 II. Activities
- III. Nucleic acids
 - A. encoding fragments, sequence, probes
 - B. mutations, chimeras, fusions
 - C. making nucleic acids
 - 10 D. vectors, cells comprising
- IV. Proteins, Peptides
 - A. fragments, sequence, immunogens, antigens
 - B. muteins
 - C. agonists/antagonists, functional equivalents
 - 15 D. making proteins
- V. Making nucleic acids, proteins
- VI. Antibodies
 - A. polyclonals
 - B. monoclonal, Kd
 - 20 C. anti-idiotypic antibodies
 - D. hybridoma cell lines
- VII. Kits and Methods to quantify IL-1 ζ
 - A. ELISA
 - B. assay mRNA encoding
 - 25 C. qualitative/quantitative
 - D. kits
- VIII. Therapeutic compositions, methods
 - A. combination compositions
 - B. unit dose
 - 30 C. administration
- IX. Receptors

I. General

Before the present compositions, formulations, and
 35 methods are described, it is to be understood that this
 invention is not limited to the particular methods,
 compositions, and cell lines described herein, as such methods,
 compositions, and cell lines may, of course, vary. It is also
 to be understood that the terminology used herein is for the
 40 purpose of describing particular embodiments only, and is not
 intended to limit the scope of the present invention which is
 only defined by the appended claims.

As used herein, including the appended claims, singular
 forms of words such as "a," "an," and "the" include their
 45 corresponding plural referents unless the context clearly
 dictates otherwise. Thus, e.g., reference to "an organism"
 includes one or more different organisms, reference to "a

cell" includes one or more of such cells, and reference to "a method" includes reference to equivalent steps and methods known to a person of ordinary skill in the art, and so forth.

Unless otherwise defined, all technical and scientific
5 terms used herein have the same meaning as commonly understood by a person of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods
10 and materials are described below. All publications, patent applications, patents, and other references discussed above are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the invention is not entitled
15 to antedate such disclosure by virtue of its prior invention. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety including all figures and drawings.

The present invention provides the amino acid sequence
20 and DNA sequence of a mammalian, e.g., human, interleukin-1 like molecule having particular defined properties, both structural and biological. This has been designated herein as interleukin-1 ζ , and increases the number of members of the IL-1 family from 6 to 7. Various cDNAs encoding these
25 molecules were obtained from primate, e.g., human, cDNA sequence libraries. Rodent counterparts should also exist. The nucleic acids encompassed herein include DNA, cDNA, and RNA sequences which encode IL-1 ζ . It is understood that nucleic acids encoding all or a portion of IL-1 ζ polypeptides
30 are also encompassed, so long as they encode a polypeptide with IL-1 ζ activity. Such nucleic acids include both naturally occurring and intentionally manipulated nucleic acids. For example, IL-1 ζ may be subjected to site-directed mutagenesis.

35 Some of the standard methods applicable are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory,

Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and periodic
5 supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York; each of which is incorporated herein by reference.

A complete nucleotide (SEQ ID NO: 1) and corresponding amino acid sequence (SEQ ID NO: 2) of a primate IL-1 ζ coding
10 segment is shown in Table 1. An alternative sequence, perhaps an allelic variant, is presented as SEQ ID NO: 3 and 4. Table 2 provides an alignment of selected family members.

Table 1: Nucleotide and amino acid sequences (see SEQ ID NO: 1-4) of mammalian, e.g., primate, IL-1 ζ . The coding sequence does not indicate a signal sequence, which has been reported for various forms of messages encoding other members of the IL-1 family. It is likely that another form of the message probably encodes a signal sequence, much like the IL-1 β prodomain which is cleaved by a convertase-like enzyme, see Dinarello (1994) FASEB J. 1314-1325).

5	CGGTTTGTTT TCTTTAGAGA TTTTACAGTG TTGGTTATAA TTGTGCACTT AATCTTTATT	60
10	TTCCTTATAC AGTAGTCCCC CCGATCAACT GGGGGCATGT TCCATACCCC TGGTGGATTC	120
	CTGAAACTGC CAGTTAGTAC CAAACCCTAT ATAGATTGTG TTTTTCCTG TACGCAGGCC	180
15	GACACACAGG AAATCATAAG TCAGGAGGGC CACTGCCACG CAGGAAAGAC CCATCTGAAC	240
	TGCTGCAAAA GCTCCGTGTC GATTTATTGC TTCCACAAAT AGTGCCGATA TGCACCAGGC	300
20	ACTGTTGTAA AACTGAAAAT ATGTTTGGG ATGTGCCCAG TCTACCTAGC TCCTTCAAGT	360
	AAAGGATCCT GAGAACTGAA GGCAAACAGA GCTCCAGGAG TCCAAGACAG AGCCACACAC	420
	CACGAGGATC CTGGCCCAGG TCTTGACTT CCATTCCCAT TTCTGTTGAG TAATAAACTC	480
25	AACGTTGAAA ATG TCC TTT GTG GGG GAG AAC TCA GGA GTG AAA ATG GGC	529
	Met Ser Phe Val Gly Glu Asn Ser Gly Val Lys Met Gly	
	1 5 10	
30	TCT GAG GAC TGG GAA AAA GAT GAA CCC CAG TGC TGC TTA GAA GAC CCG	577
	Ser Glu Asp Trp Glu Lys Asp Glu Pro Gln Cys Cys Leu Glu Asp Pro	
	15 20 25	
35	GCT GGA AGC CCC CTG GAA CCA GGC CCA AGC CTC CCC ACC ATG AAT TTT	625
	Ala Gly Ser Pro Leu Glu Pro Gly Pro Ser Leu Pro Thr Met Asn Phe	
	30 35 40 45	
40	GTT CAC ACA AGT CGA AAG GTG AAG AGC TTA AAC CCG AAG AAA TTC AGC	673
	Val His Thr Ser Arg Lys Val Lys Ser Leu Asn Pro Lys Lys Phe Ser	
	50 55 60	
45	ATT CAT GAC CAG GAT CAC AAA GTA CTG GTC CTG GAC TCT GGG AAT CTC	721
	Ile His Asp Gln Asp His Lys Val Leu Val Leu Asp Ser Gly Asn Leu	
	65 70 75	
50	ATA GCA GTT CCA GAT AAA AAC TAC ATA CGC CCA GAG ATC TTC TTT GCA	769
	Ile Ala Val Pro Asp Lys Asn Tyr Ile Arg Pro Glu Ile Phe Phe Ala	
	80 85 90	
55	TTA GCC TCA TCC TTG AGC TCA GCC TCT GCG GAG AAA GGA AGT CTG ATT	817
	Leu Ala Ser Ser Leu Ser Ser Ala Ser Ala Glu Lys Gly Ser Leu Ile	
	95 100 105	
60	CTC CTG GGG GTC TCT AAA GGG GAG TTT TGT CTC TAC TGT GAC AAG GAT	865
	Leu Leu Gly Val Ser Lys Gly Glu Phe Cys Leu Tyr Cys Asp Lys Asp	
	110 115 120 125	
	AAA GGA CAA AGT CAT CCA TCC CTT CAG CTG AAG AAG GAG AAA CTG ATG	913
	Lys Gly Gln Ser His Pro Ser Leu Gln Leu Lys Lys Glu Lys Leu Met	
	130 135 140	

	AAG CTG GCT GCC CAA AAG GAA TCA GCA CGC CGG CCC TTC ATC TTT TAT	961
	Lys Leu Ala Ala Gln Lys Glu Ser Ala Arg Arg Pro Phe Ile Phe Tyr	
	145 150 155	
5	AGG GCT CAG GTG GGC TCC CGG AAC ATG CTG GAG TCG GCG GCT CAC CCC	1009
	Arg Ala Gln Val Gly Ser Arg Asn Met Leu Glu Ser Ala Ala His Pro	
	160 165 170	
10	GGA TGG TTC ATC TGC ACC TCC TGC AAT TGT AAT GAG CCT GTT GGG GTG	1057
	Gly Trp Phe Ile Cys Thr Ser Cys Asn Cys Asn Glu Pro Val Gly Val	
	175 180 185	
15	ACA GAT AAA TTT GAG AAC AGG AAA CAC ATT GAA TTT TCA TTT CAA CCA	1105
	Thr Asp Lys Phe Glu Asn Arg Lys His Ile Glu Phe Ser Phe Gln Pro	
	190 195 200 205	
	GTT TGC AAA GCT GAA ATG AGC CCC AGT GAG GTC AGC GAT TAGGAACTG	1154
	Val Cys Lys Ala Glu Met Ser Pro Ser Glu Val Ser Asp	
	210 215	
20	CCCCATTGAA CGCCTTCCTC GCTAATTTGA ACTAATTGTA TAAAAACCCC AACCTGCTC	1214
	ACTAAAAAAA A	1225
25	MSFVGENSEGVKMGSEDWEKDEPQCCLEDPAGSPLEPGPSLPTMNFVHTSRKVKSLNPKKFSI HDQDHKVLVLDSGNLIAVPDKNYIRPEIFFALASSLSSASAEKGSLLILGVSKGEFCLYCDK DKGQSHPSLQLKKEKLMKLAAQKESARRPFIFYRAQVGSRNMLESAHPGWFICTSCNCNEP VGVTDKFENRKHIEFSFQPVCKAEMSPSEVSD	
30	Sequence of a second variant form, which exhibits nucleotide base changes at nucleotide positions of the coding region: 92 (G->T; changing amino acids G->V); 124 (A->G; changing T->A); 149 (G->C; changing R->P); 161 (G->A; changing S->N); 323 (T->C; changing L->P); and 490 (C->T; changing R->W).	
35	ATG TCC TTT GTG GGG GAG AAC TCA GGA GTG AAA ATG GGC TCT GAG GAC	48
	Met Ser Phe Val Gly Glu Asn Ser Gly Val Lys Met Gly Ser Glu Asp	
	1 5 10 15	
40	TGG GAA AAA GAT GAA CCC CAG TGC TGC TTA GAA GAC CCG GCT GTA AGC	96
	Trp Glu Lys Asp Glu Pro Gln Cys Cys Leu Glu Asp Pro Ala Val Ser	
	20 25 30	
45	CCC CTG GAA CCA GGC CCA AGC CTC CCC GCC ATG AAT TTT GTT CAC ACA	144
	Pro Leu Glu Pro Gly Pro Ser Leu Pro Ala Met Asn Phe Val His Thr	
	35 40 45	
50	AGT CCA AAG GTG AAG AAC TTA AAC CCG AAG AAA TTC AGC ATT CAT GAC	192
	Ser Pro Lys Val Lys Asn Leu Asn Pro Lys Lys Phe Ser Ile His Asp	
	50 55 60	
55	CAG GAT CAC AAA GTA CTG GTC CTG GAC TCT GGG AAT CTC ATA GCA GTT	240
	Gln Asp His Lys Val Leu Val Leu Asp Ser Gly Asn Leu Ile Ala Val	
	65 70 75 80	
	CCA GAT AAA AAC TAC ATA CGC CCA GAG ATC TTC TTT GCA TTA GCC TCA	288
	Pro Asp Lys Asn Tyr Ile Arg Pro Glu Ile Phe Phe Ala Leu Ala Ser	
	85 90 95	
60	TCC TTG AGC TCA GCC TCT GCG GAG AAA GGA AGT CCG ATT CTC CTG GGG	336

Table 2: IL-1a_human is SEQ ID NO: 5; IL-1a_mouse is SEQ ID NO: 6;
 IL-1g_human is SEQ ID NO: 7; IL-1g_mouse is SEQ ID NO: 8; IL-1b_human is
 SEQ ID NO: 9; IL-1b_mouse is SEQ ID NO: 10; IL-1x_human (IL-1RA) is SEQ
 ID NO: 11; IL-1x_mouse (IL-1RA) is SEQ ID NO: 12; IL-1d_mouse is SEQ ID
 NO: 13; IL-1e_mouse is SEQ ID NO: 14; and IL-1e_human is SEQ ID NO: 15.

	IL-1a_human	SA	PFSFLSNVKY
	IL-1a_mouse	SA	PYTYQSDLRY
10	IL-1g_human		
	IL-1g_mouse		
	IL-1b_human		
	IL-1b_mouse		
	IL-1x_humanCRPS
	IL-1x_mouseCRPS
15	IL-1d_mouse		
	IL-1z_human	MSFVGENSEGV	KMGSEDWEKD	EPQCCLEDPA	GSPLEPGPSL		PTMNFVHTSR
	IL-1e_mouseMNKEK
	IL-1e_humanMRGTP		GDADGGGRAV
20	IL-1a_human	NFMRIIKYEF	ILNDALN...	QSIIRAND..	QYLTAALHN	LD...	EAVKF
	IL-1a_mouse	KLMKLVQRKF	VMNDSLND...	QTIYQDVD.K	HYLSTTWLND	LQ...	QEVKF
	IL-1g_human	..DYFGKLES	KLSVIRNLND	QVLFIDQGNR	PLFEDMTDSD	CRDNAPRTIF	
	IL-1g_mouse	..DNFGRLLHC	TTAVIRNIND	QVLFVDKR.Q	PVFEDMTDID	QSASEPQTRL	
	IL-1b_human	.DAPVRSLNC	TLRDSQQ...	KSLVMSGP..	YELKALHLQG	QDM.EQQVVF	
25	IL-1b_mouse	.DVPIRQLHY	RLRDEQQ...	KSLVLSDP..	YELKALHLNG	QNI.NQQVIF	
	IL-1x_human	GRKSSKMQAF	RIWDVNQ...	KTFYLRN...	NQLVAGYLQG	PNV.NLEEKI	
	IL-1x_mouse	GKRPCMKQAF	RIWDTNQ...	KTFYLRN...	NQLIAGYLQG	PNI.KLEEKI	
	IL-1d_mouse	MMVLGALCF	RMKDSAL...	KVLYLHN...	NQLLAGGLHA	EKVIKGEIS	
	IL-1z_human	KVKSLNPKKF	SIHQDQDH...	KVLVLDS...	GNLIAVPDKN	YIR..PEIFF	
30	IL-1e_mouse	ELRAASPSLR	HVQDLSS...	RVWILQN...	NILTAVPRKE	QTV..PVTIT	
	IL-1e_human	YQSMCKPITG	TINDLNQ...	QVWTLQG...	QNLVAVPRSD	SVT..PVTVA	
35	IL-1a_human	DMGAYKSSK.	.DDAKITVIL	RISK.TQLYV	TAQD....ED	QPVLLKEMPE	
	IL-1a_mouse	DMYAYSSGG.	.DDSKYPVTL	KISD.SQLFV	SAQG....ED	QPVLLKELPE	
	IL-1g_human	IISMYKDS..	.QPRGMAVTI	SVKCEKISTL	SCEN.....	KIISFKEMNP	
	IL-1g_mouse	IIYMYKDS..	.EVRGLAVTL	SVKDSKMSTL	SCKN.....	KIISFEEMDP	
	IL-1b_human	SMSFVQGEE.	.SNDKIPVAL	GLKE.KNLYL	SCVL.KD.DK	PTLQLESVDP	
	IL-1b_mouse	SMSFVQGEF.	.SNDKIPVAL	GLKG.KNLYL	SCVM.KD.GT	PTLQLESVDP	
	IL-1x_human	DVVPIEP...HALFL	GIHG.GKLCL	SCVK.SG.DE	TRLQLEAVNI	
40	IL-1x_mouse	DMVPIDL...HSVFL	GKGY.YKLYM	SCVK.SG.DD	IKLQLEEVNI	
	IL-1d_mouse	VVPNRALD..	..ASLSPVIL	GVQG.GSQCL	SCGT..E.KG	PILKLEPVNI	
	IL-1z_human	ALASSLSSAS	.AEKGSLLIL	GVSK.GEFCL	YCDKDKGQSH	PSLQLKKEKL	
	IL-1e_mouse	LLPCQYLDLTL	ETNRGDPTYM	GVQR.PMSCL	FCTK..DGEQ	PVLQLGEGNI	
	IL-1e_human	VITCKYPEAL	EQGRGDPIYL	GIQN.PEMCL	YCEK..VGEQ	PTLQLKEQKI	
45	IL-1a_human	IPKTITG..S	ETNLLFFWET	HG...TKNYF	TSVAHPNLF	ATKQ...DYW	
	IL-1a_mouse	TPKLITG..S	ETDLIFFWKS	IN...SKNYF	TSAAYPELF	ATKE...QSR	
	IL-1g_human	PDNIKD...T	KSDIIFQRS	VPGHDNKMVF	ESSSYEGYFL	ACEKERDLFK	
	IL-1g_mouse	PENIDD...I	QSDLIFFQKR	VPGH.NKMEF	ESSLYEGHFL	ACQKEDDAFK	
50	IL-1b_human	KNYPKK..KM	EKRFFVFNKIE	IN...NKLEF	ESAQFPNWYI	STSQA.ENMP	
	IL-1b_mouse	KQYPKK..KM	EKRFFVFNKIE	VK...SKVEF	ESAQFPNWYI	STSQA.EHKP	
	IL-1x_human	TDLSENK.KQ	DKRFAFIRSD	SG...PTTSF	ESAACPGWFL	CTAME.ADQP	
	IL-1x_mouse	TDLSENK.EE	DKRFTFIRSE	KG...PTTSF	ESAACPGWFL	CTTLE.ADRP	
	IL-1d_mouse	MELYLGA.KE	SKSFTFYRRD	MG...LTSSF	ESAAYPGWFL	CTSPE.ADQP	
55	IL-1z_human	MKLAAQKESA	RRPFIFYRAQ	VG...SRNML	ESAHPGWFI	CTSCN.CNEP	
	IL-1e_mouse	MEMYNKK.EP	VKASLFYHKK	SG...TTSTF	ESAHPGWFI	AVCSK.GSCP	
	IL-1e_human	MDLYGQP.EP	VKPFLFYRAK	TG...RTSTL	ESVAFPDWFI	ASSKR..DQP	

Table 2 (continued):

	IL-1a_human	VCLAG.....	.GPPSITDFQ	ILENQA.....
	IL-1a_mouse	VHLAR.....	.GLPSMTDFQ	IS.....
5	IL-1g_human	LILKKEDE..	.LGDRSIMFT	VQNEDE.....
	IL-1g_mouse	LILKKKDE..	.NGDKSVMFT	LTNLHQSE...
	IL-1b_human	VFLGGTK...	.GGQDITDFT	MQFVSS.....
	IL-1b_mouse	VFLGNNS...	..GQDIIDFT	MESVSS.....
	IL-1x_human	VSLTNMPD..	.EGVMVTKFY	FQEDE.....
10	IL-1x_mouse	VSLTNTPE..	.EPLIVTKFY	FQEDQ.....
	IL-1d_mouse	VRLTQIPEDP	AWDAPITDFY	FQQCD.....
	IL-1z_human	VGVTDKFE..	..NRKHIEFS	FQPVCKAEMS	PSEVSD
	IL-1e_mouse	LILTQELG..	..EIFITDFE	MIVVH.....
15	IL-1e_human	IILTSELG..	..KSYNTAFE	LNIND.....

Table 3: Relationship between various IL-1 family members, % identity for proteins:

	hla	m1a	h1b	m1b	h1RA	m1RA	h1g	m1g	h1e	m1e	m1d	h1z
hla												
m1a	54											
h1b	21	20										
m1b	20	20	78									
h1RA	18	16	25	26								
m1RA	21	18	27	29	77							
h1g	13	16	15	16	18	17						
m1g	15	17	15	15	15	15	63					
h1e	17	16	21	18	30	29	14	12				
m1e	18	16	22	22	25	27	15	16	46			
m1d	20	19	26	27	45	45	15	13	28	28		
h1z	13	20	22	24	26	28	11	13	22	18	20	

5

Comparison of the sequences will also provide an evolutionary tree. This can be generated, e.g., using the TreeView program in combination with the ClustalX analysis software program. See Thompson, et al. Nuc. Acids Res. 25:4876-4882; and TreeView, Page, IBLIS, University of Glasgow, e-mail rpage@bio.gla.ac.uk; <http://taxonomy.zoology.gla.ac.uk.rod.treeview.html>.

10

β conformation boundaries for IL-1 ζ (SEQ ID NO: 2) are approximately: β 1 lys58-asn64; β 2 val69-ser74; β 3 asn76-val80; β 4 phe91-ser96; β 5 ser107-val113; β 6 phe118-lys126; β 7 pro131-lys136; β 8 phe154-val161; β 9 ser163-ser169; β 10 phe176-ser180; 5 β 11 glu185-gln204; and β 12 phe201-gln204. The presence of amino acid residues between β conformations β 4 and β 5 are characteristic of IL-1 agonists. IL-1 family molecules have highly conserved residues in the region encompassing β conformations β 9 and β 10.

10 Various sites for interaction with receptor are: Site A includes residues corresponding to positions of SEQ ID NO: 2 numbered 63-66, 72-74, 78, 80-87, 181-186, and 202 and 204; Site B includes residues corresponding to positions numbered 53-56, 58, 95-103, 159, 161-164, 205, and 207; and Site C 15 includes residues corresponding to positions numbered 127-153. See, e.g., USSN 09/097,976, which is incorporated herein by reference.

As used herein, the term IL-1 ζ shall be used to describe a protein comprising a protein or peptide segment having or 20 sharing the amino acid sequence shown in Table 1, or a substantial fragment thereof. The invention also includes protein variations of the IL-1 ζ allele whose sequence is provided, e.g., a mutein agonist or antagonist. Typically, such agonists or antagonists will exhibit less than about 10% 25 sequence differences, and thus will often have between 1- and 11-fold substitutions, e.g., 2-, 3-, 5-, 7-fold, and others. It also encompasses allelic and other variants, e.g., natural polymorphic variants, of the protein described. "Natural" as used herein means unmodified by artifice. Typically, it will 30 bind to its corresponding biological receptor with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. The term shall also be used herein to refer to related naturally occurring forms, 35 e.g., alleles, polymorphic variants, and metabolic variants of the mammalian protein.

This invention also encompasses proteins or peptides having substantial amino acid sequence homology with the amino acid sequence in Table 1. It will include sequence variants with relatively few substitutions, e.g., typically less than
5 about 3-5.

A substantial polypeptide "fragment", or "segment", is a stretch of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14 amino acids, more
10 often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28 amino acids, and, in particularly preferred embodiments, at
15 least about 30 or more amino acids. Sequences of segments of different proteins can be compared to one another over appropriate length stretches.

Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches, if necessary, by
20 introducing gaps as required. See, e.g., Needleham, et al., (1970) J. Mol. Biol. 48:443-453; Sankoff, et al. (1983) chapter one in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison, Addison-Wesley, Reading, MA; and software packages from
25 IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group (GCG), Madison, WI; each of which is incorporated herein by reference. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions
30 within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are intended to include natural allelic and interspecies
35 variations in the cytokine sequence. Typical homologous proteins or peptides will have from 50-100% homology (if gaps can be introduced), to 60-100% homology (if conservative

substitutions are included) with an amino acid sequence segments of Table 1. Homology measures will be at least about 70%, generally at least 76%, more generally at least 81%, often at least 85%, more often at least 88%, typically at least 90%, more typically at least 92%, usually at least 94%, more usually at least 95%, preferably at least 96%, and more preferably at least 97%, and in particularly preferred embodiments, at least 98% or more. The degree of homology will vary with the length of the compared segments.

Homologous proteins or peptides, such as the allelic variants, will share most biological activities with the embodiments described in Table 1. As used herein, the term "biological activity" is used to describe, without limitation, effects on inflammatory responses and/or innate immunity. For example, it may, like IL-1 γ , exhibit synergistic induction by splenocytes of IFN- γ in combination with IL-12 or IL-2, with or without anti-type I or anti-type II IL-1 receptor antibodies, or more structural properties as receptor binding and cross-reactivity with antibodies raised against the same or a polymorphic variant of a mammalian IL-1 ζ .

The terms ligand, agonist, antagonist, and analog of, e.g., IL-1 ζ , include molecules that modulate the characteristic cellular responses to IL-1 ζ or IL-1 ζ -like proteins, as well as molecules possessing the more standard structural binding competition features of ligand-receptor interactions, e.g., where the receptor is a natural receptor or an antibody. The cellular responses likely are mediated through binding of IL-1 ζ to cellular receptors related to, but possibly distinct from, the type I or type II IL-1 receptors.

Also, a ligand is a molecule which serves either as a natural ligand to which said receptor, or an analog thereof, binds, or a molecule which is a functional analog of the natural ligand. The functional analog may be a ligand with structural modifications, or may be a wholly unrelated molecule which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists, see, e.g., Goodman, et al. (eds. 1990) Goodman &

Gilman's: The Pharmacological Bases of Therapeutics, Pergamon Press, New York.

Rational drug design may also be based upon structural studies of the molecular shapes of a receptor or antibody and other effectors or ligands. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography, Academic Press, New York, which is hereby incorporated herein by reference.

II. Activities

The IL-1 ζ polypeptides will have a number of different biological activities, e.g., in the immune system, and will include inflammatory functions or other innate immunity responses. The IL-1 ζ polypeptides are homologous to other IL-1 proteins, but each have structural differences. For example, a human IL-1 γ gene coding sequence probably has about 70% identity with the nucleotide coding sequence of mouse IL-1 γ , and similar measures of similarity will likely apply to the IL-1 ζ . At the amino acid level, there is also likely to be about 60% identity. This level of similarity suggests that the new IL-1 ζ proteins are related to the other IL-1 α , IL-1 β , IL-1RA, IL-1 γ , IL-1 δ , and IL-1 ϵ .

The mouse IL-1 γ molecule has the ability to stimulate IFN- γ production which augments NK activity in spleen cells. See Okamura, et al. (1995) Nature 378:88-91.

The activities of the mouse IL-1 α , IL-1 β , and IL-1 γ have been compared as to their ability to induce IFN- γ , alone or in combination with IL-2 or IL-12 in SCID splenocytes and purified NK cells. See Hunter, et al. (1995) J. Immunol. 155:4347-4354; and Bancroft, et al. (1991) Immunol. Revs.

124:5-24. The IL-1 γ was found to be much more potent in stimulating IFN-1 γ than either IL-1 α or IL-1 β . IL-1 ζ and agonists or antagonists should have related activities to these or to the other new IL-1 δ and IL-1 ϵ , typically affecting
5 similar immune functions, including inflammatory responses.

In IL-2 activated NK cells, IFN- γ production is blocked by the addition of anti-IL-1 β antibodies. See Hunter, et al. (1995). However, mouse IL-1 γ can overcome this block and induce IFN- γ . This is the only cytokine known to be able to do
10 this. In addition, in vivo, administration of mouse IL-1 γ to mice infected with the parasite T. Cruzi significantly decreases parasitemia.

The present disclosure also describes new assays for activities predicted for the IL-1 ζ molecules. Corresponding
15 activities should be found in other mammalian systems, including primates or rodents. It is likely that the new primate IL-1-like molecules produced by similar recombinant means to the human IL-1 γ protein should exhibit a biological activity of modulating lymphocyte cells in production of IFN- γ .
20 See assays described, e.g., in de Waal Malefyt, et al., in de Vries and de Waal Malefyt (eds. 1995) "Interleukin-10" Landes Co., Austin, TX. Furthermore, there is substantial likelihood of synergy with other IL-1 or IL-12 related agonists or antagonists. It is likely that the receptors, which are
25 expected to include multiple different polypeptide chains, exhibit species specificity for their corresponding ligands. The IL-1 α and IL-1 β ligands both signal through heterodimeric receptors.

30 III. Nucleic Acids

This invention contemplates use of isolated nucleic acid or fragments, e.g., which encode this or a closely related protein, or fragments thereof, e.g., to encode a biologically active corresponding polypeptide. The term "isolated nucleic
35 acid or fragments" as used herein means a nucleic acid, e.g., a DNA or RNA molecule, that is not immediately contiguous when present in the naturally occurring genome of the organism from

which it is derived. Thus, the term describes, e.g., a nucleic acid that is incorporated into a vector, such as a plasmid or viral vector; a nucleic acid that is incorporated into the genome of a heterologous cell (or the genome of homologous cell, but at a site different from that at which it normally occurs); and a nucleic acid that exists as a separate molecule, e.g., a DNA fragment produced by PCR amplification or restriction enzyme digestion, or an RNA molecule produced by in vitro transcription. The term also describes a recombinant (i.e., genetically engineered) nucleic acid that forms part of a hybrid gene encoding additional polypeptide sequences that can be used, e.g., in the production of a fusion protein. In addition, this invention embodies virtually any engineered or nucleic acid molecule created by artifice that encodes a biologically active protein or polypeptide having characteristic IL-1 ζ activity.

Typically, the nucleic acid is capable of hybridizing, under appropriate conditions, with a nucleic acid sequence segment shown in Table 1. Said biologically active protein or polypeptide can be a full length protein, or fragment, and will typically have a segment of amino acid sequence highly homologous to one shown in Table 1. Further, this invention covers the use of isolated or recombinant nucleic acid, or fragments thereof, which encode proteins having fragments which are homologous to the newly disclosed IL-1-like proteins. The isolated nucleic acids can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others from the natural gene.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially pure, e.g., separated from other components which naturally accompany a native sequence, such as ribosomes, polymerases, and flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates, which are thereby

distinguishable from naturally occurring compositions, and chemically synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule, either
5 completely or substantially pure.

An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some embodiments, contain heterogeneity, preferably minor. This heterogeneity is typically found at the polymer ends or portions not
10 critical to a desired biological function or activity.

A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g.,
15 involving human intervention in the nucleotide sequence. Typically this intervention involves in vitro manipulation, although under certain circumstances it may involve more classical animal breeding techniques. Alternatively, it can be a nucleic acid made by generating a sequence comprising
20 fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants as found in their natural state. Thus, e.g., products made by transforming cells with an unnaturally occurring vector is encompassed, as are nucleic
25 acids comprising sequence derived using a synthetic oligonucleotide process. Such a process is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a restriction enzyme sequence recognition site.
30 Alternatively, the process is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms, e.g., encoding a fusion protein. Restriction enzyme
35 recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences,

control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. This will include a dimeric repeat. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode similar polypeptides to fragments of the IL-1 ζ and fusions of sequences from various different interleukin or related molecules, e.g., growth factors.

A "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least 21 nucleotides, more generally at least 25 nucleotides, ordinarily at least 30 nucleotides, more ordinarily at least 35 nucleotides, often at least 39 nucleotides, more often at least 45 nucleotides, typically at least 50 nucleotides, more typically at least 55 nucleotides, usually at least 60 nucleotides, more usually at least 66 nucleotides, preferably at least 72 nucleotides, more preferably at least 79 nucleotides, and in particularly preferred embodiments will be at least 85 or more nucleotides including, e.g., 100, 150, 200, 250, etc. Preferred embodiments will exhibit a plurality of distinct, e.g., nonoverlapping, segments of the specified length. Typically, the plurality will be at least two, more usually at least three, and preferably 5, 7, or even more. While the length minima are provided, longer lengths, of various sizes, may be appropriate, e.g., one of length 7, and two of length 12. Typically, fragments of different genetic sequences can be compared to one another over appropriate length stretches, particularly defined segments such as the domains described below.

A nucleic acid which codes for an IL-1 ζ will be particularly useful to identify genes, mRNA, and cDNA species which code for itself or closely related proteins, as well as DNAs which code for polymorphic, allelic, or other genetic variants, e.g., from different individuals or related species. Preferred probes for such screens are those regions of the interleukin which are conserved between different polymorphic variants or which contain nucleotides which lack specificity,

and will preferably be full length or nearly so. In other situations, polymorphic variant specific sequences will be more useful.

This invention further covers recombinant nucleic acid molecules and fragments having a nucleic acid sequence identical to or highly homologous to the isolated DNA set forth herein. In particular, the sequences will often be operably linked to DNA segments which control transcription, translation, and DNA replication. These additional segments typically assist in expression of the desired nucleic acid segment.

Homologous nucleic acid sequences, when compared to one another or Table 1 sequences, exhibit significant similarity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. Comparative hybridization conditions are described in greater detail below.

Substantial identity in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 60% of the nucleotides, generally at least 66%, ordinarily at least 71%, often at least 76%, more often at least 80%, usually at least 84%, more usually at least 88%, typically at least 91%, more typically at least about 93%, preferably at least about 95%, more preferably at least about 96 to 98% or more, and in particular embodiments, as high at about 99% or more of the nucleotides, including, e.g., segments encoding structural domains such as the segments described below. Alternatively, substantial identity will exist when the segments will hybridize under selective hybridization conditions, to a strand or its complement, typically using a sequence derived from Table 1. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 14 nucleotides, more typically at least about 65%, preferably at

least about 75%, and more preferably at least about 90%. See, Kanehisa (1984) Nuc. Acids Res. 12:203-213. The length of homology comparison, as described, may be over longer stretches, and in certain embodiments will be over a stretch
5 of at least about 17 nucleotides, generally at least about 20 nucleotides, ordinarily at least about 24 nucleotides, usually at least about 28 nucleotides, typically at least about 32 nucleotides, more typically at least about 40 nucleotides, preferably at least about 50 nucleotides, and more preferably
10 at least about 75 to 100 or more nucleotides.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates
15 are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

20 Optical alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math. 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443, by the search for similarity method of Pearson
25 and Lipman (1988) Proc. Nat'l Acad. Sci. USA 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally Ausubel et al.,
30 supra).

One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a
35 tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng and Doolittle (1987)

J. Mol. Evol. 35:351-360. The method used is similar to the method described by Higgins and Sharp (1989) CABIOS 5:151-153. The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. For example, a reference sequence can be compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps.

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described Altschul, et al. (1990) J. Mol. Biol. 215:403-410. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul, et al., supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension of the word hits in each direction are halted when: the cumulative alignment

score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The
5 BLAST algorithm parameters W , T , and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a wordlength (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) Proc. Nat'l Acad. Sci. USA 89:10915) alignments (B) of 50, expectation (E) of
10 10, $M=5$, $N=4$, and a comparison of both strands.

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin and Altschul (1993) Proc. Nat'l Acad. Sci. USA 90:5873-5787). One
15 measure of similarity provided by the BLAST algorithm is the smallest sum probability ($P(N)$), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if
20 the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

A further indication that two nucleic acid sequences of
25 polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second
30 polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions, as described below.

35 Stringent conditions, in referring to homology in the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters

typically controlled in hybridization reactions. Stringent temperature conditions will usually include temperatures in excess of about 30° C, more usually in excess of about 37° C, typically in excess of about 45° C, more typically in excess of about 55° C, preferably in excess of about 65° C, and more preferably in excess of about 70° C. Stringent salt conditions will ordinarily be less than about 500 mM, usually less than about 400 mM, more usually less than about 300 mM, typically less than about 200 mM, preferably less than about 100 mM, and more preferably less than about 80 mM, even down to less than about 20 mM. However, the combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370, which is hereby incorporated herein by reference. Hybridization under stringent conditions should give a background of at least 2-fold over background, preferably at least 3-5 or more.

The isolated DNA can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode this protein or its derivatives. These modified sequences can be used to produce mutant proteins (muteins) or to enhance the expression of variant species. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant IL-1-like derivatives include predetermined or site-specific mutations of the protein or its fragments, including silent mutations using genetic code degeneracy. "Mutant IL-1 ζ " as used herein encompasses a polypeptide otherwise falling within the homology definition of the IL-1 ζ as set forth above, but having an amino acid sequence which differs from that of other IL-1-like proteins as found in nature, whether by way of deletion, substitution, or insertion. In particular, "site specific mutant IL-1 ζ " encompasses a protein having substantial homology with a protein of Table 1, and typically

shares most of the biological activities of the form disclosed herein.

Although site specific mutation sites are predetermined, mutants need not be site specific. Mammalian IL-1 ζ mutagenesis can be achieved by making amino acid insertions or deletions in the gene, coupled with expression. Substitutions, deletions, insertions, or combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy- terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mammalian IL-1 ζ mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis. See also Sambrook, et al. (1989) and Ausubel, et al. (1987 and periodic Supplements).

The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Polymerase chain reaction (PCR) techniques can often be applied in mutagenesis. Alternatively, mutagenesis primers are commonly used methods for generating defined mutations at predetermined sites. See, e.g., Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach and Dveksler (1995; eds.) PCR Primer: A Laboratory Manual Cold Spring Harbor Press, CSH, NY.

IV. Proteins, Peptides

As described above, the present invention encompasses mammalian IL-1 ζ , e.g., whose sequences are disclosed in Table 1, and described above. Allelic and other variants are also contemplated, including, e.g., fusion proteins combining portions of such sequences with others, including epitope tags and functional domains.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these rodent proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. Thus, the fusion product of a growth factor with an interleukin is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences.

In addition, new constructs may be made from combining similar functional or structural domains from other related proteins, e.g., growth factors or other cytokines. For example, receptor-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992, each of which is incorporated herein by reference. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of receptor-binding specificities. For example, the receptor binding domains from other related ligand molecules may be added or substituted for other domains of this or related proteins. The resulting protein will often have hybrid function and properties. For example, a fusion protein may include a targeting domain which may serve to provide sequestering of the fusion protein to a particular organ, e.g., a ligand portions which is specifically bound by spleen cells and would serve to accumulate in the spleen.

Candidate fusion partners and sequences can be selected from various sequence data bases, e.g., GenBank, c/o NCBI; and BCG, University of Wisconsin Biotechnology Computing Group, Madison, WI, which are each incorporated herein by reference.

5 The present invention particularly provides muteins which act as agonists or antagonists of the IL-1 ζ . Structural alignment of primate IL-1 ζ and other members of the IL-1 family show conserved features/residues, particularly 12 β strands folded into a β -trefoil fold. The 12 IL-1 ζ β strand domains are recited, respectively, about: β 1 lys58-asp64; β 2 val69-ser74; β 3 asn76-val80; β 4 phe91-ser96; β 5 ser107-val113; β 6 phe118-lys126; β 7 pro131-lys136; β 8 phe154-val161; β 9 ser163-ser169; β 10 phe176-ser180; β 11 glu185-gln204; and β 12 phe201-gln204. The presence of amino acid residues between β conformations β 4 and β 5 are characteristic of IL-1 agonists. IL-1 family molecules have highly conserved residues in the region encompassing β conformations β 9 and β 10.

Alignment of the primate IL-1 ζ with other members of the IL-1 family indicates that the β conformations correspond to similar sequences in other IL-1 family members. See also, Bazan, et al. (1996) Nature 379:591; Lodi, et al. (1994) Science 263:1762-1766; Sayle and Milner-White (1995) TIBS 20:374-376; and Gronenberg, et al. (1991) Protein Engineering 4:263-269.

25 The IL-1 α and IL-1 β ligands bind an IL-1 receptor type I as the primary receptor and this complex then forms a high affinity receptor complex with the IL-1 receptor type III. Such receptor subunits are probably shared with the new IL-1 family members.

30 The mouse IL-1 γ does not bind to the known mouse IL-1 receptor types I, II (decoy receptor), or III. In addition, the mouse IGIF biological activity cannot be blocked with anti-type I, II, or III antibodies. This suggests that the related mouse IGIF binds to receptors related to the IL-1 receptors already isolated, but not yet identified as
35 receptors for the IGIF.

The solved structures for IL-1 β , the natural IL-1 receptor antagonist (IL-1Ra), and a co-structure of IL-1Ra/IL-1 receptor type I, however, suggest how to make a primate antagonist for IL-1 ζ (see, e.g., accession numbers: U65590, gbU19844, gbU19845, gi2173679, gi2170133, gi2172939, gbM15300, gbM28983, gbU65590, gbM74294, embX04964, gi2169698, gi2169368 emb270047, gi914939, gi220782, embX52731, embX56972 and embX12497, for various species examples of IL-1 family members). Structural analyses of the mature primate IL-1 ζ suggest that its β -trefoil structures contact the IL-1 receptor over three binding sites (designated A, B and C). Sites A and C bind to the first receptor subunit (alpha) of IL-1 while site B binds the IL-1 second receptor subunit (beta). Homology sequence comparison of the IL-1 family members reveals that the only known antagonist to IL-1 receptor (IL-1x, or IL-1RA; Table 2) is missing an amino acid domain bounded by the β 4 and β 5 strands. This domain maps to a portion of site B in primate IL-1 ζ (Table 2) that binds to the IL-1 second receptor subunit, suggesting that its absence confers antagonist activity as evidenced by homology comparison among other IL-1 family members. This loop portion of contact site B spans approximately 7-10 amino residues, while in IL-1RA the loop is "cut off" with only 2 residues remaining. Therefore, IL-1RA binds normally to receptor type I, but cannot interact with receptor type III. This makes IL-1RA into an effective IL-1 antagonist.

The corresponding location in primate IL-1 ζ (between β 4 and β 5) defines a domain that forms a polypeptide loop which is part of a primary binding segment to the IL-1 receptor type. The loop is defined, for IL-1 ζ , approximately by amino residues ser100-gly106 of SEQ ID NO: 2. Accordingly, IL-1 ζ antagonist activity should be generated by removal all or an appropriate portion of a corresponding portion of amino acids located between β 4 and β 5. This suggests that analogous modifications to the loop between the β 4 and the β 5 strands will lead to variants with predictable biological activities. With mouse IL-1RA, it was shown that replacement of the mouse

IL-1RA residues with those mouse IL-1 β residues introduced IL-1 activity to the IL-1RA variant (IL-1RA could then bind type III receptor). Similar substitutions should establish that type III receptor can probably be used by primate IL-1 ζ or
5 muteins.

Sites A and C should mediate binding of IL-1 ζ to the first IL-1 receptor subunit, e.g., an alpha receptor subunit. Site A contacts correspond in IL-1 ζ to amino residues corresponding to positions of SEQ ID NO: 2 numbered about 63-
10 66, 72-74, 78, 80-87, 181-186, and 202 and 204; Site B includes residues corresponding to positions numbered about 53-56, 58, 95-103, 159, 161-164, 205, and 207; and Site C includes residues corresponding to positions numbered about 127-153. See, e.g., USSN 09/097,976, which is incorporated
15 herein by reference.

Similar variations in other species counterparts of IL-1 ζ ligand sequence, e.g., in the corresponding regions, should provide similar interactions with receptor. Substitutions with either mouse sequences or human sequences are indicated.
20 Conversely, conservative substitutions away from the receptor binding interaction regions will probably preserve most biological activities.

"Derivatives" of the mammalian IL-1 ζ include amino acid sequence mutants, glycosylation variants, metabolic
25 derivatives and covalent or aggregative conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionality's to groups which are found in the IL-1 ζ amino acid side chains or at the N- or C- termini, e.g., by means which are well known in the art. These derivatives
30 can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group containing residues, e.g., lysine or
35 arginine. Acyl groups are selected from the group of alkyl-moieties including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for
5 accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence
10 which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the interleukin or fragments thereof with other proteins of
15 polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred derivatization sites with cross-linking agents are at free
20 amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between the interleukin and other homologous or heterologous proteins are also provided. Homologous polypeptides may be fusions between different growth factors, resulting in, for instance, a hybrid protein
25 exhibiting ligand specificity for multiple different receptors, or a ligand which may have broadened or weakened specificity of binding to its receptor. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative
30 proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a receptor, e.g., a ligand-binding segment, so that the presence or location of a desired ligand may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609, which is
35 hereby incorporated herein by reference. Other gene fusion partners include glutathione-S-transferase (GST), bacterial β -galactosidase, trpE, Protein A, β -lactamase, alpha amylase,

alcohol dehydrogenase, and yeast alpha mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce
5 suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

10 Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments, the modifications will
15 be useful labeling reagents, or serve as purification targets, e.g., affinity ligands.

Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and
20 expression are described generally, e.g., in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1-3, Cold Spring Harbor Laboratory, and Ausubel, et al. (eds. 1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York, which are each incorporated
25 herein by reference. Techniques for synthesis of polypeptides are described, e.g., in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; each of which is
30 incorporated herein by reference. See also, Dawson, et al. (1994) Science 266:776-779 for methods to make larger polypeptides.

In another embodiment, the present invention relates to substantially purified peptide fragments of IL-1 ζ that block
35 binding between IL-1 family members and a target receptor. Such peptide fragments could represent research and diagnostic tools in the study of inflammatory reactions to antigenic

challenge and the development of more effective anti-inflammatory therapeutics. In addition, pharmaceutical compositions comprising isolated and purified peptide fragments of IL-1 ζ may represent effective anti-inflammatory therapeutics.

The term "substantially purified" as used herein refers to a molecule, such as a peptide that is substantially free of other proteins, lipids, carbohydrates, nucleic acids, or other biological materials with which it is naturally associated.

For example, a substantially pure molecule, such as a polypeptide, can be at least 60%, by dry weight, the molecule of interest. One skilled in the art can purify IL-1 ζ peptides using standard protein purification methods and the purity of the polypeptides can be determined using standard methods including, e.g., polyacrylamide gel electrophoresis (e.g., SDS-PAGE), column chromatography (e.g., high performance liquid chromatography (HPLC)), and amino-terminal amino acid sequence analysis.

The invention relates not only to fragments of naturally-occurring IL-1 ζ , but also to IL-1 ζ mutants and chemically synthesized derivatives of IL-1 ζ that block binding between IL-1 family members and a target receptor.

For example, changes in the amino acid sequence of IL-1 ζ are contemplated in the present invention. IL-1 ζ can be altered by changing the nucleic acid sequence encoding the protein. Preferably, only conservative amino acid alterations are undertaken, using amino acids that have the same or similar properties. Illustrative amino acid substitutions include the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine, glutamine, or glutamate; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine;

threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; valine to isoleucine or leucine.

Additionally, other variants and fragments of IL-1 ζ can be used in the present invention. Variants include analogs, homologues, derivatives, muteins, and mimetics of IL-1 ζ that retain the ability to block binding between IL-1 family members and a target receptor. Fragments of the IL-1 ζ refer to portions of the amino acid sequence of IL-1 ζ as defined in SEQ ID NO: 2 that also retain this ability. The variants and fragments can be generated directly from IL-1 ζ itself by chemical modification, by proteolytic enzyme digestion, or by combinations thereof. Additionally, genetic engineering techniques, as well as methods of synthesizing polypeptides directly from amino acid residues, can be employed.

Non-peptide compounds that mimic the binding and function of IL-1 ζ ("mimetics") can be produced by the approach outlined in Saragovi, et al. (1991) Science 253:792-795. Mimetics are molecules which mimic elements of protein secondary structure. See, e.g., Johnson et al., "Peptide Turn Mimetics," in Pezzuto, et al. (eds. 1993) Biotechnology and Pharmacy, Chapman and Hall, New York. The underlying rationale behind the use of peptide mimetics is that the peptide backbone of proteins exists chiefly to orient amino acid side chains in such a way as to facilitate molecular interactions. For the purposes of the present invention, appropriate mimetics can be considered to be the equivalent of IL-1 ζ .

Variants and fragments also can be created by recombinant techniques employing genomic or cDNA cloning methods. Site-specific and region-directed mutagenesis techniques can be employed. See, e.g., vol. 1, ch. 8 in Ausubel, et al. (eds. 1989 and periodic updates) Current Protocols in Molecular Biology Wiley and Sons; and Oxender and Fox (eds.) Protein Engineering Liss, Inc. In addition, linker-scanning and PCR-mediated techniques can be employed for mutagenesis. See, e.g., Erlich (ed. 1989) PCR Technology Stockton Press. Protein sequencing, structure and modeling approaches for use with the above techniques are disclosed, e.g., in Oxender and

Fox (eds.) Protein Engineering Liss, Inc; and Ausubel, et al. (eds. 1989 and periodic updates) Current Protocols in Molecular Biology Wiley and Sons.

5 This invention also contemplates the use of derivatives of IL-1 ζ other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and
10 (3) adsorption complexes, e.g., with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of a receptor or other binding molecule, e.g., an antibody. For example, an IL-1 ζ ligand can
15 be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated SEPHAROSE, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of IL-1 ζ receptor, antibodies, or other
20 similar molecules. The IL-1 ζ can also be labeled with a detectable group, e.g., radio-iodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays.

25 An IL-1 ζ of this invention can be used as an immunogen for the production of antisera or antibodies specific, e.g., capable of distinguishing between other IL-1 family members and an IL-1 ζ , for the interleukin or fragments thereof. The purified interleukin can be used to screen monoclonal
30 antibodies or antigen-binding fragments prepared by immunization with various forms of impure preparations containing the protein. In particular, the term "antibodies" also encompasses antigen binding fragments of natural antibodies. The purified interleukin can also be used as a
35 reagent to detect antibodies generated in response to the presence of elevated levels of expression, or immunological disorders which lead to antibody production to the endogenous

cytokine. Additionally, IL-1 ζ fragments may also serve as immunogens to produce the antibodies of the present invention, as described immediately below. For example, this invention contemplates antibodies having binding affinity to or being
5 raised against the amino acid sequence shown in Table 1, fragments thereof, or homologous peptides. In particular, this invention contemplates antibodies having binding affinity to, or having been raised against, specific fragments which are predicted to be, or actually are, exposed at the exterior
10 protein surface of the native cytokine.

The blocking of physiological response to these interleukins may result from the inhibition of binding of the ligand to the receptor, likely through competitive inhibition. Thus, in vitro assays of the present invention will often use
15 antibodies or ligand binding segments of these antibodies, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either binding region mutations and modifications, or ligand mutations and modifications, e.g., ligand analogs.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to the interleukin or fragments compete with a test compound for binding to a receptor or antibody. In this manner, the neutralizing antibodies or fragments can be used to detect the
20 presence of a polypeptide which shares one or more binding sites to a receptor and can also be used to occupy binding sites on a receptor that might otherwise bind an interleukin.
25

V. Making Nucleic Acids and Protein

DNA which encodes the protein or fragments thereof can be
30 obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples. Natural sequences can be isolated using standard methods and the sequences provided
35 herein, e.g., in Table 1. Other species counterparts can be identified by hybridization techniques, or by various PCR

techniques, combined with or by searching in sequence databases.

This DNA can be expressed in a wide variety of host cells for the synthesis of a full-length interleukin or fragments which can in turn, e.g., be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified agonist/antagonist molecules; and for structure/function studies. Each variant or its fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The protein, or portions thereof, may be expressed as fusions with other proteins.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired receptor gene or its fragments, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control elements are capable of effecting expression within a suitable host. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

The vectors of this invention include those which contain DNA which encodes a protein, as described, or a fragment thereof encoding a biologically active equivalent polypeptide. The DNA can be under the control of a viral promoter and can

encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNA coding for such a protein in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNA coding for the receptor is inserted into the vector such that growth of the host containing the vector expresses the cDNA in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the interleukin protein or its fragments in various hosts using vectors that do not contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of the protein encoding portion or its fragments into the host DNA by recombination.

Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into the genome of the host. Expression vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids are the most commonly used form of vector but all other forms of vectors which serve an equivalent function and which are, or become, known in the art are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory Manual, Elsevier, N.Y., and Rodriguez, et al. (eds.) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, 1988, which are incorporated herein by reference.

Transformed cells are cells, preferably mammalian, that have been transformed or transfected with receptor vectors constructed using recombinant DNA techniques. Transformed host cells usually express the desired protein or its fragments, but for purposes of cloning, amplifying, and manipulating its DNA, do not need to express the subject

protein. This invention further contemplates culturing transformed cells in a nutrient medium, thus permitting the interleukin to accumulate in the culture. The protein can be recovered, either from the culture or from the culture medium.

5 For purposes of this invention, nucleic sequences are operably linked when they are functionally related to each other. For example, DNA for a pre-sequence or secretory leader is operably linked to a polypeptide if it is expressed as a pre-protein or participates in directing the polypeptide
10 to the cell membrane or in secretion of the polypeptide. A promoter is operably linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to permit translation. Usually, operably linked
15 means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression.

Suitable host cells include prokaryotes, lower
20 eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., E. coli and B. subtilis. Lower eukaryotes include yeasts, e.g., S. cerevisiae and Pichia, and species of the genus Dictyostelium. Higher eukaryotes include established tissue culture cell
25 lines from animal cells, both of non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used herein, E. coli
30 and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express the receptor or its fragments include, but are not limited to, such vectors as
35 those containing the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540).

See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-, and Ipp-derived Promoters", in Vectors: A Survey of Molecular Cloning Vectors and Their Uses, (eds. Rodriguez and Denhardt), Butterworth, Boston, Chapter 10, pp. 205-236, which is incorporated herein by reference.

Lower eukaryotes, e.g., yeasts and Dictyostelium, may be transformed with IL-1 ζ sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, Saccharomyces cerevisiae. It will be used to generically represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the receptor or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YE ρ -series); integrating types (such as the YIp-series), or mini-chromosomes (such as the YCp-series).

Higher eukaryotic tissue culture cells are normally the preferred host cells for expression of the functionally active interleukin protein. In principle, virtually any higher eukaryotic tissue culture cell line is workable, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred. Transformation or transfection and propagation of such cells has become a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if

genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from
5 adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo PolyA, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such
10 as pAC 373 or pAC 610.

For secreted proteins, an open reading frame usually encodes a polypeptide that consists of a mature or secreted product covalently linked at its N-terminus to a signal
15 peptide. The signal peptide is cleaved prior to secretion of the mature, or active, polypeptide. The cleavage site can be predicted with a high degree of accuracy from empirical rules, e.g., von-Heijne (1986) Nucleic Acids Research 14:4683-4690, and the precise amino acid composition of the signal peptide
20 does not appear to be critical to its function, e.g., Randall, et al. (1989) Science 243:1156-1159; Kaiser et al. (1987) Science 235:312-317.

It will often be desired to express these polypeptides in a system which provides a specific or defined glycosylation
25 pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For
30 example, the interleukin gene may be co-transformed with one or more genes encoding mammalian or other glycosylating enzymes. Using this approach, certain mammalian glycosylation patterns will be achievable in prokaryote or other cells.

The source of IL-1 ζ can be a eukaryotic or prokaryotic
35 host expressing recombinant IL-1 ζ DNA, such as is described above. The source can also be a cell line such as mouse Swiss 3T3 fibroblasts, but other mammalian cell lines are also

contemplated by this invention, with the preferred cell line being from the human species.

Now that the entire sequence is known, the priamte IL-1 ζ , fragments, or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis, Springer-Verlag, New York; and Bodanszky (1984) The Principles of Peptide Synthesis, Springer-Verlag, New York; all of each which are incorporated herein by reference. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (e.g., p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the foregoing processes.

The IL-1 ζ protein, fragments, or derivatives are suitably prepared in accordance with the above processes as typically employed in peptide synthesis, generally either by a so-called stepwise process which comprises condensing an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to the terminal amino acid. Amino groups that are not being used in the coupling reaction typically must be protected to prevent coupling at an incorrect location.

If a solid phase synthesis is adopted, the C-terminal amino acid is bound to an insoluble carrier or support through its carboxyl group. The insoluble carrier is not particularly limited as long as it has a binding capability to a reactive carboxyl group. Examples of such insoluble carriers include halomethyl resins, such as chloromethyl resin or bromomethyl resin, hydroxymethyl resins, phenol resins, tert-alkyloxycarbonylhydrazidated resins, and the like.

An amino group-protected amino acid is bound in sequence through condensation of its activated carboxyl group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After
5 synthesizing the complete sequence, the peptide is split off from the insoluble carrier to produce the peptide. This solid-phase approach is generally described by Merrifield, et al. (1963) in J. Am. Chem. Soc. 85:2149-2156, which is incorporated herein by reference.

10 The prepared protein and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, e.g., by extraction, precipitation, electrophoresis, various forms of chromatography, and the like. The interleukin of this invention can be obtained in
15 varying degrees of purity depending upon its desired use. Purification can be accomplished by use of the protein purification techniques disclosed herein, see below, or by the use of the antibodies herein described in methods of immunoabsorbant affinity chromatography. This immunoabsorbant
20 affinity chromatography is carried out by first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate cells, lysates of other cells expressing the interleukin, or lysates or supernatants of cells producing the protein as a result of
25 DNA techniques, see below.

Generally, the purified protein will be at least about 40% pure, ordinarily at least about 50% pure, usually at least about 60% pure, typically at least about 70% pure, more typically at least about 80% pure, preferable at least about
30 90% pure and more preferably at least about 95% pure, and in particular embodiments, 97%-99% or more. Purity will usually be on a weight basis, but can also be on a molar basis. Different assays will be applied as appropriate.

35 VI. Antibodies

The term "antibody" or "antibody molecule" as used in this invention includes intact molecules as well as fragments

thereof, such as Fab, $F(ab')_2$, and Fv which are capable of binding the epitopic determinant. These antibody fragments retain some ability to selectively bind with its antigen or receptor and are defined as follows: (1) Fab, the fragment
5 which contains a monovalent antigen-binding fragment of an antibody molecule can be produced by digestion of whole IgG antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain; (2) Fab', the fragment of an antibody molecule can be obtained by treating whole IgG
10 antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two Fab' fragments are obtained per antibody molecule; (3) $(Fab')_2$, the fragment of the IgG antibody that can be obtained by treating whole antibody with the enzyme pepsin without subsequent
15 reduction; $F(ab')_2$ is a dimer of two Fab' fragments held together by two disulfide bonds; (4) Fv, defined as a genetically engineered fragment containing essentially the variable region of the light chain and the variable region of the heavy chain expressed as two chains; and (5) Single Chain
20 Antibody ("SCA"), defined as a genetically engineered molecule containing essentially the variable region of the light chain, the variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule.

25 Methods of making these fragments are known in the art. See, e.g., Harlow and Lane (current edition) Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, New York. Therefore, the phrase "antibody molecule" in its various forms as used herein contemplates both an intact antibody
30 (immunoglobulin) molecule and an immunologically active portion of an antibody (immunoglobulin) molecule. Recombinant methods may be applied to make these fragments.

The term "monoclonal antibody" refers to a population of one species of antibody molecule of antigen-specificity. A
35 monoclonal antibody contains one species of antibody combining site capable of immunoreacting with a particular antigen and thus typically displays a single binding affinity for that

antigen. A monoclonal antibody may therefore contain a bispecific antibody molecule having two antibody combining sites, each immunospecific for a different antigen. In one embodiment, the first antibody molecule is affixed to a solid support. In addition, the antibody molecules in a phage display combinatorial library are also monoclonal antibodies.

As used in this invention, the term "epitope" means an antigenic determinant on an antigen to which the paratope of an antibody binds. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics.

The word "complex" as used herein refers to the product of a specific binding agent-ligand reaction. An exemplary complex is an immunoreaction product formed by an antibody-antigen reaction.

The term "antigen" refers to a polypeptide or protein that is able to selectively bind to (immunoreact with) an antibody and form an immunoreaction product (immunocomplex). The site on the antigen to which the antibody binds is referred to as an antigenic determinant or epitope, and the labeling should be detectable, e.g., 2X, 5X, or more above background.

The method of the invention for detection of antibodies that bind to novel epitopes in a sample is performed in vitro, e.g., in immunoassays in which the antibodies can be identified in liquid phase or bound to a solid phase carrier. Preferably, the method is performed with a capture antibody bound to a solid support. Preferably, the capture antibody is a monoclonal antibody molecule.

Examples of types of immunoassays which can be utilized to detect novel antibodies in a sample, include competitive and non-competitive immunoassays in either a direct or indirect format. Examples of such immunoassays are the radioimmunoassay (RIA) and the sandwich (immunometric) assay. Detection of the antibodies can be done utilizing immunoassays

which are run in either the forward, reverse, or simultaneous modes, including competition immunoassays and immunohistochemical assays on physiological samples.

Preferably, the method of the invention utilizes a forward immunoassay. Those of skill in the art will know, or can readily discern, other immunoassay formats without undue experimentation.

Solid phase-bound antibody molecules are bound by adsorption from an aqueous medium, although other modes of affixation, such as covalent coupling or other well known means of affixation to the solid matrix can be used. Preferably, the first antibody molecule is bound to a support before forming an immunocomplex with antigen, however, the immunocomplex can be formed prior to binding the complex to the solid support.

Non-specific protein binding sites on the surface of the solid phase support are preferably blocked. After adsorption of solid phase-bound antibodies, an aqueous solution of a protein free from interference with the assay such as bovine, horse, or other serum albumin that is also free from contamination with the antigen is admixed with the solid phase to adsorb the admixed protein onto the surface of the antibody-containing solid support at protein binding sites on the surface that are not occupied by the antibody molecule.

A typical aqueous protein solution contains about 2-10 weight percent bovine serum albumin in PBS at a pH of about 7-8. The aqueous protein solution-solid support mixture is typically maintained for a time period of at least one hour at a temperature of about 4°-37° C and the resulting solid phase is thereafter rinsed free of unbound protein.

The first preselected antibody can be bound to many different carriers and used to detect novel epitope binding antibodies in a sample. Examples of well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amyloses, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the carrier can be either soluble or insoluble for purposes of the

invention. Those skilled in the art will know of other suitable carriers for binding antibodies, or will be able to ascertain such, using routine experimentation.

In addition, if desirable, an antibody for detection in these immunoassays can be detectably labeled in various ways. There are many different labels and methods of labeling known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include enzymes, radioisotopes, fluorescent compounds, colloidal metals, chemiluminescent compounds, and bio-luminescent compounds. Those of ordinary skill in the art will know of other suitable labels for binding to the monoclonal antibodies of the invention, or will be able to ascertain such, using routine experimentation. Furthermore, the binding of these labels to the antibodies used in the method of the invention can be done using standard techniques common to those of ordinary skill in the art.

Antibodies which bind to IL-1 ζ polypeptides of the invention can be prepared using an intact polypeptide or fragments containing small peptides of interest as the immunizing antigen. The polypeptide or a peptide used to immunize an animal can be derived from translated cDNA or chemical synthesis which can be conjugated to a carrier protein, if desired. Such commonly used carriers which are chemically coupled to the peptide include keyhole limpet hemocyanin (KLH), thyroglobulin, bovine serum albumin (BSA), and tetanus toxoid. The coupled peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit).

If desired, polyclonal or monoclonal antibodies can be further purified, e.g., by binding to and elution from a matrix to which the polypeptide or a peptide to which the antibodies were raised is bound. Those of skill in the art will know of various techniques common in the immunology arts for purification and/or concentration of polyclonal antibodies, as well as monoclonal antibodies. See, e.g., Coligan, et al. (current ed.) Current Protocols in Immunology, Wiley Interscience.

It is also possible to use the anti-idiotypic technology to produce monoclonal antibodies which mimic an epitope. For example, an anti-idiotypic monoclonal antibody made to a first monoclonal antibody will have a binding domain in the
5 hypervariable region which is the "image" of the epitope bound by the first monoclonal antibody.

The preparation of polyclonal antibodies is well-known to those skilled in the art. See, e.g., Green, et al. "Production of Polyclonal Antisera" pages 1-5 in Manson (ed.)
10 Immunochemical Protocols Humana Press; Harlow and Lane; and Coligan, et al. Current Protocols in Immunology.

The preparation of monoclonal antibodies likewise is conventional. See, e.g., Kohler and Milstein (1975) Nature 256:495-497; Coligan, et al., sections 2.5.1-2.6.7; and Harlow
15 and Lane Antibodies: A Laboratory Manual, Cold Spring Harbor Press. Briefly, monoclonal antibodies can be obtained by injecting mice with a composition comprising an antigen, verifying the presence of antibody production by removing a serum sample, removing the spleen to obtain B lymphocytes,
20 fusing the B lymphocytes with myeloma cells to produce hybridomas, cloning the hybridomas, selecting positive clones that produce antibodies to the antigen, and isolating the antibodies from the hybridoma cultures. Monoclonal antibodies can be isolated and purified from hybridoma cultures by a
25 variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A Sepharose, size-exclusion chromatography, and ion-exchange chromatography. See, e.g., Coligan, et al.; Barnes, et al. "Purification of Immunoglobulin G (IgG)" in Methods in
30 Molecular Biology, vol. 10, pages 79-104 (Humana Press, current ed.). Methods of in vitro and in vivo multiplication of monoclonal antibodies are well-known to those skilled in the art. Multiplication in vitro may be carried out in suitable culture media such as Dulbecco's Modified Eagle
35 Medium or RPMI 1640 medium, optionally replenished, e.g., by a mammalian serum such as fetal calf serum or trace elements and growth-sustaining supplements such as normal mouse peritoneal

exudate cells, spleen cells, bone marrow macrophages. Production in vitro provides relatively pure antibody preparations and allows scale-up to yield large amounts of the desired antibodies. Large scale hybridoma cultivation can be carried out by homogenous suspension culture in an airlift reactor, in a continuous stirrer reactor, or in immobilized or entrapped cell culture. Multiplication in vivo may be carried out by injecting cell clones into mammals histocompatible with the parent cells, e.g., syngeneic mice, to cause growth of antibody-producing tumors. Optionally, the animals are primed with a hydrocarbon, especially oils such as pristane (tetramethylpentadecane) prior to injection. After one to three weeks, the desired monoclonal antibody is recovered from the body fluid of the animal.

Therapeutic applications are conceivable for the antibodies of the present invention. For example, antibodies of the present invention may also be derived from subhuman primate antibody. General techniques for raising therapeutically useful antibodies in baboons may be found, e.g., in Goldenberg, et al. (1991) WO 91/11465; and Losman, et al. (1990) Int. J. Cancer 46:310-314.

Alternatively, a therapeutically useful anti-IL-1 ζ antibody may be derived from a "humanized" monoclonal antibody. Humanized monoclonal antibodies are produced by transferring mouse complementary determining regions from heavy and light variable chains of the mouse immunoglobulin into a human variable domain, and then substituting human residues in the framework regions of the murine counterparts. The use of antibody components derived from humanized monoclonal antibodies obviates potential problems associated with the immunogenicity of murine constant regions. General techniques for cloning murine immunoglobulin variable domains are described, e.g., by Orlandi, et al. (1989) Proc. Nat'l Acad. Sci. USA 86:3833-3837. Techniques for producing humanized monoclonal antibodies are described, e.g., by Jones et al. (1986) Nature 321:522-525; Riechmann, et al. (1988) Nature 332:323-327; Verhoeyen, et al. (1988) Science 239:1534-

1536; Carter, et al. (1992) Proc. Nat'l Acad. Sci. USA
89:4285-4289; Sandhu (1992) Crit. Rev. Biotech. 12:437-462;
and Singer, et al. (1993) J. Immunol. 150:2844-2857.

Antibodies of the invention also may be derived from
5 human antibody fragments isolated from a combinatorial
immunoglobulin library. See, e.g., Barbas, et al. (1991)
Methods: A Companion to Methods in Enzymology, vol. 2, page
119; and Winter, et al. (1994) Ann. Rev. Immunol. 12:433-455.
Cloning and expression vectors that are useful for producing a
10 human immunoglobulin phage library can be obtained, e.g., from
STRATAGENE Cloning Systems (La Jolla, CA).

In addition, antibodies of the present invention may be
derived from a human monoclonal antibody. Such antibodies are
obtained from transgenic mice that have been "engineered" to
15 produce specific human antibodies in response to antigenic
challenge. In this technique, elements of the human heavy and
light chain loci are introduced into strains of mice derived
from embryonic stem cell lines that contain targeted
disruptions of the endogenous heavy and light chain loci. The
20 transgenic mice can synthesize human antibodies specific for
human antigens, and the mice can be used to produce human
antibody-secreting hybridomas. Methods for obtaining human
antibodies from transgenic mice are described by Green, et al.
(1994) Nature Genet. 7:13; Lonberg, et al. (1994) Nature
25 368:856; and Taylor, et al. (1994) Int. Immunol. 6:579.

Antibody fragments of the present invention can be
prepared by proteolytic hydrolysis of the antibody or by
expression in *E. coli* of DNA encoding the fragment. Antibody
fragments can be obtained by pepsin or papain digestion of
30 whole antibodies by conventional methods. For example,
antibody fragments can be produced by enzymatic cleavage of
IgG antibodies with pepsin to provide a 5S fragment denoted
F(ab')₂. This fragment can be further cleaved using a thiol
reducing agent, and optionally a blocking group for the
35 sulfhydryl groups resulting from cleavage of disulfide
linkages, to produce 3.5 S Fab' monovalent fragments.
Alternatively, an enzymatic cleavage using papain produces two

monovalent Fab fragments and an Fc fragment directly. These methods are described, e.g., by Goldenberg, U.S. patents No. 4,036,945 and No. 4,331,647, and references contained therein. These patents are hereby incorporated in their entireties by
5 reference including all figures, drawings, and illustrations. See also Nisonhoff, et al. (1960) Arch. Biochem. Biophys. 89:230-xxx; Porter (1959) Biochem. J. 73:119-xxx; Edelman, et al. (1967) Methods in Enzymology, vol. 1, Academic Press; and Coligan, et al.

10 Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical, or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the
15 intact antibody.

For example, Fv fragments comprise an association of V_H and V_L chains. This association may be noncovalent, as described in Inbar, et al. (1972) Proc. Nat'l Acad. Sci. USA 69:2659. Alternatively, the variable chains can be linked by
20 an intermolecular disulfide bond or cross-linked by chemicals such as glutaraldehyde. See, e.g., Sandhu (1992) Crit. Rev. Biotech. 12:437. Preferably, the Fv fragments comprise V_H and V_L chains connected by a peptide linker. These single-chain antigen binding proteins (sFv) are prepared by constructing a
25 structural gene comprising DNA sequences encoding the V_H and V_L domains connected by an oligonucleotide. The structural gene is inserted into an expression vector, which is subsequently introduced into a host cell such as E. coli. The recombinant host cells synthesize a single polypeptide chain
30 with a linker peptide bridging the two V domains. Methods for producing sFvs are described, e.g., by Whitlow, et al. (1991) Methods: A Companion to Methods in Enzymology, vol. 2, page 97; Bird, et al. (1988) Science 242:423-426; Ladner, et al., U.S. patent No. 4,946,778; Pack, et al. (1993) Bio/Technology
35 11:1271-77; and Sandhu (1992) Crit. Rev. Biotech. 12:437-462.

Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR

peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, e.g., by using the polymerase chain reaction to synthesize the variable region
5 from RNA of antibody-producing cells. See, e.g., Larrick, et al. (1991) Methods: A Companion to Methods in Enzymology, vol. 2, page 106.

Antibodies can be raised to the various mammalian, e.g., primate IL-1 ζ polypeptides, both in naturally occurring native
10 forms and in their denatured forms, the difference being that antibodies to the active ligand are more likely to recognize epitopes which are only present in the native conformations. Denatured antigen detection can also be useful in, e.g., Western analysis. Anti-idiotypic antibodies are also
15 contemplated, which would be useful as agonists or antagonists of a natural receptor or an antibody.

A number of immunogens may be used to produce antibodies selectively reactive with IL-1 ζ proteins. Recombinant protein is the preferred immunogen for the production of monoclonal or
20 polyclonal antibodies. Naturally occurring protein may also be used either in pure or impure form. Synthetic peptides made using the IL-1 ζ protein sequences described herein may also be used as an immunogen for the production of antibodies to the antigens. Recombinant protein can be expressed in
25 eukaryotic or prokaryotic cells as described herein, and purified as described. The product is then injected into an animal capable of producing antibodies. Either monoclonal or polyclonal antibodies may be generated for subsequent use in immunoassays to measure the protein.

30 Methods of producing polyclonal antibodies are known to those of skill in the art. In brief, an immunogen, preferably a purified protein, is mixed with an adjuvant and animals are immunized with the mixture. The animal's immune response to the immunogen preparation is monitored by taking test bleeds
35 and determining the titer of reactivity to the protein antigen of interest. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal

and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired. See Harlow and Lane.

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell. Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according to the general protocol outlined by Huse, et al. (1989) Science 246:1275-1281.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of the protein can be raised by immunization of animals with conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective protein, or screened for agonistic or antagonistic activity. These monoclonal antibodies will usually bind with at least a K_D of about 1 mM, more usually at least about 300 μ M, typically at least about 100 μ M, more typically at least about 30 μ M, preferably at least about 10 μ M, and more preferably at least about 3 μ M or better; including 1 μ M, 300 nM, 100 nM, 30 nM, etc.

The antibodies, including antigen binding fragments, of this invention can have significant diagnostic or therapeutic value. They can be potent antagonists that bind to the interleukin and inhibit binding to the receptor or inhibit the ability of IL-1 ζ to elicit a biological response. They also

can be useful as non-neutralizing antibodies and can be coupled to toxins or radionuclides to bind producing cells, or cells localized to the source of the interleukin. Further, these antibodies can be conjugated to drugs or other
5 therapeutic agents, either directly or indirectly by means of a linker.

The antibodies of this invention can also be useful in diagnostic applications. As capture or non-neutralizing antibodies, they can bind to the interleukin without
10 inhibiting receptor binding. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying IL-1 ζ . They may be used as reagents for Western blot analysis, or for immunoprecipitation or immunopurification of the respective
15 protein.

Protein fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. Mammalian IL-1 ζ and its fragments may be fused or covalently linked to a variety
20 of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York; and Williams, et al. (1967) Methods in Immunology and
25 Immunochemistry, Vol. 1, Academic Press, New York; each of which are incorporated herein by reference, for descriptions of methods of preparing polyclonal antisera. A typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the
30 repeated immunizations and the gamma globulin is isolated.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g.,
35 Stites, et al. (eds.) Basic and Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A

Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York; and particularly in Kohler and Milstein (1975) in Nature 256: 495-497, which discusses one method of generating
5 monoclonal antibodies. Each of these references is incorporated herein by reference. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a
10 hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of
15 immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to
20 selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-546, each of which is hereby incorporated
25 herein by reference. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a
30 detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic
35 particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241.

Also, recombinant or chimeric immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; or made in transgenic mice, see Mendez, et al. (1997) Nature Genetics 15:146-156. These references are incorporated herein by reference.

5 The antibodies of this invention can also be used for affinity chromatography in isolating the IL-1 ζ . Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, SEPHADEX, or the like, where a cell lysate may be passed through the column,
10 the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified protein will be released. The protein may be used to purify antibody.

 The antibodies may also be used to screen expression libraries for particular expression products. Usually the
15 antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

 Antibodies raised against an IL-1 ζ will also be used to raise anti-idiotypic antibodies. These will be useful in
20 detecting or diagnosing various immunological conditions related to expression of the protein or cells which express receptors for the protein. They also will be useful as agonists or antagonists of the interleukin, which may be competitive inhibitors or substitutes for naturally occurring
25 ligands.

Binding Agent:IL-1 ζ Polypeptide Complex

 An IL-1 ζ polypeptide that specifically binds to or that is specifically immunoreactive with an antibody, e.g., such as
30 a polyclonal antibody, generated against a defined immunogen, e.g., such as an immunogen consisting of an amino acid sequence of SEQ ID NO: 2 or fragments thereof or a polypeptide generated from the nucleic acid of SEQ ID NO: 1 is typically determined in an immunoassay. Included within the metes and
35 bounds of the present invention are those nucleic acid sequences described herein, including functional variants, that encode polypeptides that selectively bind to polyclonal

antibodies generated against the prototypical IL-1 ζ polypeptide as structurally and functionally defined herein. The immunoassay typically uses a polyclonal antiserum which was raised, e.g., to a protein of SEQ ID NO: 2. This
5 antiserum is selected to have low crossreactivity against other IL-1 family members, preferably from the same species, and such crossreactivity is removed by immunoabsorption or depletion prior to use in the immunoassay. Appropriate selective serum preparations can be isolated, and
10 characterized.

In order to produce antisera for use in an immunoassay, the protein of SEQ ID NO: 2 is isolated as described herein. For example, recombinant protein may be produced in a mammalian cell line. An appropriate host, e.g., an inbred
15 strain of mice such as Balb/c, is immunized with the protein of SEQ ID NO: 2 using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane). Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a
20 carrier protein can be used as an immunogen. Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, e.g., a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of 10^4 or greater are selected and tested for
25 their cross reactivity against other IL-1 family members, e.g., IL-1 α , IL-1 β , IL-1RA, IL-1 γ , IL-1 δ , and IL-1 ϵ , using a competitive binding immunoassay such as the one described in Harlow and Lane, supra, at pages 570-573. Preferably at least two IL-1 family members are used in this determination in
30 conjunction with IL-1 ζ . These IL-1 family members can be produced as recombinant proteins and isolated using standard molecular biology and protein chemistry techniques as described herein. Thus, antibody preparations can be identified or produced having desired selectivity or
35 specificity for subsets of IL-1 family members.

Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For example, the

protein of SEQ ID NO: 2 can be immobilized to a solid support. Proteins added to the assay compete with the binding of the antisera to the immobilized antigen. The ability of the above proteins to compete with the binding of the antisera to the
5 immobilized protein is compared to the protein of SEQ ID NO: 2. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% crossreactivity with each of the proteins listed above are selected and pooled. The cross-reacting antibodies
10 are then removed from the pooled antisera by immunoabsorption with the above-listed proteins.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein to the immunogen protein. In order to make
15 this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required is less than twice the amount of the
20 protein of SEQ ID NO: 2 that is required, then the second protein is said to specifically bind to an antibody generated to the immunogen.

It is understood that these IL-1 ζ polypeptide is a member of a family of homologous proteins that comprise at least 6 so
25 far identified genes. For a particular gene product, such as the IL-1 ζ , the term refers not only to the amino acid sequences disclosed herein, but also to other proteins that are allelic, non-allelic or species variants. It also understood that the term "IL-1 ζ " includes nonnatural mutations
30 introduced by deliberate mutation using conventional recombinant technology such as single site mutation, or by excising short sections of DNA encoding the respective proteins, or by substituting new amino acids, or adding new amino acids. Such minor alterations must substantially
35 maintain the immunoidentity of the original molecule and/or its biological activity. Thus, these alterations include proteins that are specifically immunoreactive with a

designated naturally occurring IL-1 related protein, e.g., the IL-1 ζ polypeptide shown in SEQ ID NO: 2. The biological properties of the altered proteins can be determined by expressing the protein in an appropriate cell line and measuring the appropriate effect upon lymphocytes. Particular protein modifications considered minor would include conservative substitution of amino acids with similar chemical properties, as described above for the IL-1 family as a whole. By aligning a protein optimally with the protein of SEQ ID NO: 2 and by using the conventional immunoassays described herein to determine immunoidentity, one can determine the protein compositions of the invention.

VII. Kits and quantitation

Both naturally occurring and recombinant forms of the IL-1 like molecules of this invention are particularly useful in kits and assay methods. For example, these methods would also be applied to screening for binding activity, e.g., receptors for these proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds per year. See, e.g., a BIOMEK automated workstation, Beckman Instruments, Palo Alto, California, and Fodor, et al. (1991) Science 251:767-773, which is incorporated herein by reference. The latter describes means for testing binding by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays to screen for a receptor or agonist/antagonist homologous proteins can be greatly facilitated by the availability of large amounts of purified, soluble IL-1 ζ in an active state such as is provided by this invention.

Purified IL-1 ζ can be coated directly onto plates for use in the aforementioned receptor screening techniques. However, non-neutralizing antibodies to these proteins can be used as capture antibodies to immobilize the respective interleukin on the solid phase, useful, e.g., in diagnostic uses.

This invention also contemplates use of IL-1 ζ polypeptides and their fusion products in a variety of

diagnostic kits and methods for detecting the presence of the protein or its receptor. Alternatively, or additionally, antibodies against the molecules may be incorporated into the kits and methods. Typically the kit will have a compartment
5 containing either a defined IL-1 ζ peptide or gene segment or a reagent which recognizes one or the other. Typically, recognition reagents, in the case of peptide, would be a receptor or antibody, or in the case of a gene segment, would usually be a hybridization probe.

10 A preferred kit for determining the concentration of, e.g., IL-1 ζ , a sample would typically comprise a labeled compound, e.g., receptor or antibody, having known binding affinity for IL-1 ζ , a source of IL-1 ζ (naturally occurring or recombinant) as a positive control, and a means for separating
15 the bound from free labeled compound, e.g., a solid phase for immobilizing the IL-1 ζ in the test sample. Compartments containing reagents, and instructions, will normally be provided.

Antibodies, including antigen binding fragments, specific
20 for mammalian IL-1 ζ or a peptide fragment, or receptor fragments are useful in diagnostic applications to detect the presence of elevated levels of IL-1 ζ and/or its fragments. Diagnostic assays may be homogeneous (without a separation step between free reagent and antibody-antigen complex) or
25 heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA) and the
30 like. For example, unlabeled antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to IL-1 ζ or to a particular fragment thereof. These assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A
35 Laboratory Manual, CSH., and Coligan (ed. 1991 and periodic supplements) Current Protocols In Immunology Greene/Wiley, New York.

Anti-idiotypic antibodies may have similar use to serve as agonists or antagonists of IL-1 ζ . These should be useful as therapeutic reagents under appropriate circumstances.

Frequently, the reagents for diagnostic assays are
5 supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody, or labeled receptor is provided. This is usually in conjunction with other additives, such as buffers,
10 stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent, and will contain instructions for
15 proper use and disposal of reagents. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium having appropriate concentrations for performing the assay.

The aforementioned constituents of the diagnostic assays
20 may be used without modification or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In these assays, a test compound, IL-1 ζ , or antibodies thereto can be labeled
25 either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as ^{125}I , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence
30 intensity, wavelength shift, or fluorescence polarization. Both of the patents are incorporated herein by reference. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

35 There are also numerous methods of separating the bound from the free ligand, or alternatively the bound from the free test compound. The IL-1 ζ can be immobilized on various

matrixes followed by washing. Suitable matrices include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the receptor to a matrix include, without limitation, direct adhesion to plastic, use of a capture
5 antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of antibody/antigen complex by appropriate methods including those utilizing, e.g., an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable
10 separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30:1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678, each of which is incorporated herein by
15 reference.

The methods for linking protein or fragments to various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through
20 the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

25 Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequence of an IL-1 ζ . These sequences can be used as probes for detecting levels of the IL-1 ζ in patients suspected of having an immunological disorder. The preparation of both RNA
30 and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least about 18 nucleotides, and the
35 polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly ^{32}P . However, other techniques may also be

employed, such as using biotin modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected. The use of probes to the novel anti-sense RNA may be carried out in conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART). This also includes amplification techniques such as polymerase chain reaction (PCR).

Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97.

VIII. Therapeutic Utility

This invention provides reagents with significant therapeutic value. The IL-1 ζ polypeptides (naturally occurring or recombinant), mutein agonists and antagonists, and antibodies, along with compounds identified as having binding affinity to the interleukin or its receptor or antibodies, should be useful in the treatment of conditions exhibiting abnormal expression of the interleukin. Such abnormality will typically be manifested by immunological disorders. Additionally, this invention should provide therapeutic value in various diseases or disorders associated with abnormal expression or abnormal triggering of response to the interleukin. The mouse IL- γ has been suggested to be

involved in tumors, allergies, and infectious diseases, e.g., pulmonary tuberculosis, leprosy, fulminant hepatitis, and viral infections, such as HIV. The IL-1 ζ or antagonist may have similar function, suggesting combination compositions
5 with other agonists or antagonists of IL-1 family members.

T helper cells mediate effector functions in infectious, allergic, or autoimmune diseases through production of cytokines. CD4 positive T cells can be divided into Th1 and Th2 subsets on the basis of their cytokine profile upon
10 antigen stimulation. Recently obtained evidence has shown that Th1 and Th2 cells differ in responsiveness and receptor expression for IL-1 family molecules. See, e.g., Robinson, et al. (1997) Immunity 7:571-581. Whereas Th1 cells respond to IL-1 γ , Th2 cells respond to IL-1 α . This differential
15 responsiveness between Th1 and Th2 cells to IL-1 γ and IL-1 α , respectively, may have profound implications for regulation of ongoing Th cell responses. The novel IL-1 molecules described here could play a similar role in either supporting a Th1 or Th2 response, depending on the presence or absence of their
20 cognate IL-1 receptors on the cell surface of these immune cells; e.g., IL-1RD4 (ST2) is an orphan IL-1-like receptor exclusively expressed on the Th2 subset. See, e.g., Lohning, et al. (1998) Proc. Nat'l Acad. Sci. USA 95:6930-6935; and USSN 09/040,714, which are incorporated herein by reference.

25 In addition, the dendritic cell expression profile shows human IL-1 γ primarily expressed in activated dendritic cells. Activated dendritic cells are also a major producer of IL-12, and it is thought that this dendritic cell produced IL-12 plays a major role in directing a Th1 type response. The
30 combination of IL-1 γ and IL-12 should be extremely potent in inducing IFN- γ , suggesting that IL-1 ζ , or antagonists thereof, may have similar function. It is possible that the combination of pro-inflammatory cytokines under certain circumstances could lead to septic shock. An antagonist,
35 mutein or antibody, could prove very useful in this situation. See Rich (ed.) Clinical Immunology: Principles and Practice, Mosby.

Additionally, IL-1 ζ being homologous members of the IL-1 family likely play a role in modulating of local and systemic inflammatory processes (see, Durum, et al. (1986) Ann. Rev. Immunol. 3:253-xxx), through the enhancement of blood flow, induction of chemoattractants, and the enhancement and adherence of inflammatory cells such as macrophages and neutrophils at the site of inflammation. Additionally, it is possible that IL-1 ζ can induce fibroblast growth and may play a role in contributing to the pathogenesis of chronic inflammation, as in rheumatoid arthritis or periodontal disease.

IL-1 ζ is also likely to play a role in systemic inflammatory reactions, such as fever, hypoglycemia, the acute phase response of the liver, reduced plasma iron and zinc, and increased plasma copper. A systemic reaction such as septic shock involves vasodilation, due to IL-1, most likely in combination with other cytokines, including, e.g., TNF, IFN- γ , and leukemia inhibitory factor (LIF). The newly described IL-1 ζ is also likely to be similarly involved.

Recombinant IL-1 ζ , mutein agonists or antagonists, or IL-1 ζ antibodies can be purified and then administered to a patient. These reagents can be combined for therapeutic use with additional active ingredients, e.g., in conventional pharmaceutically innocuous stabilizers and excipients. These combinations can be sterilized, e.g., in conventional dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments thereof which are not complement binding.

Receptor screening using IL-1 ζ or fragments thereof can be performed to identify molecules having binding affinity to the interleukin. Subsequent biological assays can then be utilized to determine if a receptor can provide competitive binding, which can block intrinsic stimulating activity. Receptor fragments can be used as a blocker or antagonist in that it blocks the activity of IL-1 ζ . Likewise, a compound

having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of IL-1 ζ . This invention further contemplates the therapeutic use of antibodies to IL-1 ζ as antagonists.

5 The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and
10 efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Various considerations
15 are described, e.g., in Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences (current ed.), Mack Publishing Co., Easton, Penn.; each of which is hereby incorporated herein by reference. Methods for
20 administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the
25 Merck Index, Merck & Co., Rahway, New Jersey. Because of the likely high affinity binding between an IL-1 ζ and its receptors, low dosages of these reagents would be initially expected to be effective. And the signaling pathway suggests extremely low amounts of ligand may have effect. Thus, dosage
30 ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10 μ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier.
35 Slow release formulations, or slow release apparatus will often be utilized for continuous administration.

IL-1 ζ polypeptides, and antibodies or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in a conventional dosage formulation. While it is possible for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier must be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; Avis, et al. (eds. 1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, NY; Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Tablets Dekker, NY; and Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, NY.

Another therapeutic approach included within the invention involves direct administration of reagents or compositions by conventional administration techniques (e.g., but not restricted to local injection, inhalation, or administered systemically), to the subject with an inflammatory disorder. The reagent, formulation or composition may also be targeted to specific cells or receptors by, e.g., methods described herein. The actual dosage of reagent, formulation or composition that modulates

an inflammatory disorder depends on many factors, including the size and health of an organism, however one of one of ordinary skill in the art can use the following teachings describing the methods and techniques for determining clinical dosages. See, e.g., Spilker (1984) Guide to Clinical Studies and Developing Protocols, Raven Press, New York; Spilker (1991) Guide to Clinical Trials, Raven Press, New York; Craig and Stitzel (eds. 1986) Modern Pharmacology 2d ed., Little, Brown, Boston; Speight (ed. 1987) Avery's Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics, 3d ed., Williams and Wilkins, Baltimore; and Tallarida, et al. (1988) Principles in General Pharmacology, Springer-Verlag, New York; which describe how to determine the appropriate dosage; but, generally, in the range of about between 0.5 ng/ml and 500 µg/ml inclusive final concentration are administered per day to an adult in a pharmaceutically-acceptable carrier. The therapy of this invention may be combined with or used in association with other therapeutic agents, particularly agonists or antagonists of other IL-1 family members.

IX. Receptors

The description of the IL-1ζ ligand herein provides means to identify a receptor, as described above. Such receptor should bind specifically to the IL-1ζ with reasonably high affinity. Various constructs are made available which allow either labeling of the IL-1ζ to detect its receptor. For example, directly labeling IL-1ζ, fusing onto it markers for secondary labeling, e.g., FLAG or other epitope tags, etc., will allow detection of receptor. This can be histological, as an affinity method for biochemical purification, or labeling or selection in an expression cloning approach. A two-hybrid selection system may also be applied making appropriate constructs with the available IL-1ζ sequences. See, e.g., Fields and Song (1989) Nature 340:245-246. Typically, a cytokine will bind to its receptor at a Kd of at least about 30 µM, preferably at least about 10 µM, and more

preferably at least about 3 μM or better; including 1 μM , 300 nM, 100 nM, 30 nM, etc.

EXAMPLES

I. General Methods

Some of the standard methods are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and Supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York. Methods for protein purification include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Deutscher (1990) "Guide to Protein Purification" in Meth. Enzymol., vol. 182, and other volumes in this series; and manufacturer's literature on use of protein purification products, e.g., Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA. Combination with recombinant techniques allow fusion to appropriate segments, e.g., to a FLAG sequence or an equivalent which can be fused via a protease-removable sequence. See, e.g., Hochuli (1989) Chemische Industrie 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering, Principle and Methods 12:87-98, Plenum Press, N.Y.; and Crowe, et al. (1992) QIAexpress: The High Level Expression & Protein Purification System QIAGEN, Inc., Chatsworth, CA.

Computer sequence analysis is performed, e.g., using available software programs, including those from the GCG (U. Wisconsin) and GenBank sources. Public sequence databases were also used, e.g., from GenBank and others.

Many techniques applicable to IL-4, IL-10, and IL-1 γ , IL-1 δ , and IL-1 ϵ may be applied to IL-1 ζ , as described, e.g., in U.S. Patent No. 5,017,691 (IL-4), USSN 07/453,951 (IL-10), USSN 08/110,683 (IL-10 receptor); USSN 08/651,998 (IL-1 γ); USSN

09/062,866 or USSN 09/097,976 (IL-1 δ and IL-1 ϵ), each of which is incorporated herein by reference for all purposes.

II. Amplification of IL-1 ζ fragment by PCR

5 There are various methods of isolating the DNA sequences encoding IL-1 ζ polypeptides. For example, DNA is isolated from a genomic or cDNA library using labeled oligonucleotide probes having sequences identical or complementary to the sequences disclosed herein. Full-length probes may be used,
10 or oligonucleotide probes may be generated by comparison of the sequences disclosed. Such probes can be used directly in hybridization assays to isolate DNA encoding IL-1 ζ polypeptides, or probes can be designed for use in amplification techniques such as PCR, for the isolation of DNA
15 encoding IL-1 ζ polypeptides.

 Various methods of amplifying target sequences, such as the polymerase chain reaction, can also be used to prepare DNA encoding IL-1 ζ polypeptides. Polymerase chain reaction (PCR) technology is used to amplify such nucleic acid sequences
20 directly from mRNA, from cDNA, and from genomic libraries or cDNA libraries. The isolated sequences encoding IL-1 ζ polypeptides may also be used as templates for PCR amplification.

 In PCR techniques, oligonucleotide primers complementary
25 to two 5' regions in the DNA region to be amplified are synthesized. The polymerase chain reaction is then carried out using the two primers. See Innis, et al. (current eds.) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA. Primers can be selected to amplify the
30 entire regions encoding a full-length IL-1 ζ polypeptide or to amplify smaller DNA segments, as desired. Once such regions are PCR-amplified, they can be sequenced and oligonucleotide probes can be prepared from sequence obtained using standard techniques. These probes can then be used to isolate DNA's
35 encoding IL-1 ζ polypeptides.

 Oligonucleotides for use as probes are chemically synthesized, e.g., according to the solid phase

phosphoramidite triester method first described by Beaucage and Carruthers (1983) Tetrahedron Lett. 22:1859-1862, or using an automated synthesizer, as described in Needham-VanDevanter, et al. (1984) Nucleic Acids Res. 12:6159-6168. Purification
5 of oligonucleotides is performed, e.g., by native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson and Regnier (1983) J. Chrom. 255: 137-149. The sequence of the synthetic oligonucleotide can be verified using the chemical degradation method of Maxam and Gilbert in
10 Grossman and Moldave (eds. 1980) Methods in Enzymology 65: 499-560 Academic Press, New York.

The peptide segments, along with comparison to homologous genes, can also be used to produce appropriate oligonucleotides to screen a library. The genetic code can be
15 used to select appropriate oligonucleotides useful as probes for screening. In combination with polymerase chain reaction (PCR) techniques, synthetic oligonucleotides will be useful in selecting desired clones from a library.

Complementary sequences will also be used as probes or
20 primers. Based upon identification of the likely amino terminus, other peptides should be particularly useful, e.g., coupled with anchored vector or poly-A complementary PCR techniques or with complementary DNA of other peptides.

To identify a homologous IL-1 ζ polypeptide, degenerate
25 oligonucleotides are designed which corresponded to conserved regions among known IL-1 family members. The primers are used for polymerase chain reactions on primate genomic DNA followed by subcloning the PCR products using restriction sites placed at the 5' ends of the primers, picking individual E. coli
30 colonies carrying these subcloned inserts, and using a combination of random sequencing and hybridization analysis to eliminate known IL-1 family members.

Subsequently, PCR products are gel-purified, digested with appropriate restriction enzymes, gel-purified again, and
35 subcloned in the Bluescript vector (Stratagene, San Diego, CA). Bacterial colonies carrying individual subclones are picked into 96 well microtiter plates, and multiple replicas

are prepared by plating the cells onto nitrocellulose. The replicate filters are hybridized to probes representing known members of the IL-1 family, and DNA is prepared from non-hybridizing colonies for sequence analysis.

- 5 Two appropriate forward and reverse primers are selected using the sequences supplied herein (see Table 1) and common knowledge. See, e.g., Innis, et al. (current eds.) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach and Dveksler (current eds.) PCR
10 Primer: A Laboratory Manual Cold Spring Harbor Press, CSH, NY. RT-PCR is used on an appropriate mRNA sample selected for the presence of message to produce a cDNA, e.g., a monocyte or macrophage cell sample.

- Full length clones may be isolated by hybridization of
15 cDNA libraries from appropriate tissues pre-selected by PCR signal.

- As is commonly known, PCR primers are typically designed to contain at least 15 nucleotides, e.g., 15-30 nucleotides. The design of IL-1 ζ specific primers containing 21
20 nucleotides, e.g., that code for IL-1 ζ polypeptides containing at least 6 amino acids from the IL-1 ζ domains are described as follows. Other PCR primers designed to amplify other IL-1 ζ polypeptide fragments are designed in a similar fashion, e.g., mutagenesis primers. Preferably, most or all of the
25 nucleotides in such a primer encode conserved amino acids, e.g., amino residues of SEQ. ID NO: 2 or 4, including IL-1 ζ -specific amino acids as described herein. For example, primers containing at least 40% IL-1 ζ -conserved amino acids can be used. Such a primer, containing 21 nucleotides, can
30 include sequences encoding at least 3/7, 4/7, 5/7, 6/7 or 7/7 IL-1 ζ -conserved amino acids. Once IL-1 ζ amino acids are selected as templates against which primer sequences are to be designed, the primers can be synthesized using, e.g., standard chemical methods. Due to the degeneracy of the genetic code
35 and the bias of preferred species variants, such primers should be designed to include appropriate degenerate

sequences, as can be readily determined using common knowledge.

Based on the guidelines presented above, IL-1 ζ -conserved amino acid peptides can be used as templates for the design of IL-1 ζ specific primers. Additional examples can be found by analysis of sequence alignments of IL-1 ζ polypeptides (Table 1 and 2). Primers can be designed to amplify various structural features or domains, e.g., a 4-10 amino acid region of either IL-1 ζ polypeptide that corresponds to one of the 12 β strands could be amplified using this strategy. Depending on the length of the primer, desired primers can be designed, e.g., to correspond to 4-7 consecutive amino acids of the segments. Preferred segments include, e.g., GENSGVK; EDWEKD; CCLEDPA; FVHTSR; KKFSIHD; VLVLDS; NLIAVP; FFALAS; SSASAEK; SLILLGV; FCLYCDK; PSLQLK; KLMKLAAQ; FIFYRAQ; SRNMLES; WFICTS; EPVGVGT; or FSFQPVC (see SEQ ID NO: 2); or FVHTSP; SPILLGV; or SWNMLES (see SEQ ID NO: 4). Longer preferred segments include, e.g., GVKMGSEDWEKD; AGSPLEPGPSLP; SRKVKSLNPKKF; HDQDHKVLVLDS; NLIAVPDKNYIR; FALASSLSSASA; GQSHPSLQLKKE; MKLAAQKESARR; FYRAQVGSRNML; TSCNCNEPVGVT; FENRKHIEFSFQ; or PVCKAEMSPSEV (see SEQ ID NO: 2); or AVSPLEPGPSLP; SPKVKNLNPCKF; or FYRAQVGSWNML (see SEQ ID NO: 4).

As is described above, IL-1 ζ primers, e.g., primers based on IL-1 ζ specific peptides shown above, or portions thereof, can be used in PCR reactions to generate IL-1 ζ probes which can be used in standard screening methods to identify nucleic acids encoding IL-1 family members (see e.g., Ausubel, et al., supra).

III. Tissue distribution of IL-1 ζ

Message for the gene encoding IL-1 ζ has been detected in a mixture of fetal lung, testis, and B cell cDNAs. PCR has provided positive signals in cDNA libraries derived from normal colon, lung, and a marginally detectable signal from ovary. It was detectable by Northern blotting in a polyA+ sample from a colorectal adenocarcinoma.

Southern Analysis: DNA (5 µg) from a primary amplified cDNA library is digested with appropriate restriction enzymes to release the inserts, run on a 1% agarose gel and transferred to a nylon membrane (Schleicher and Schuell, Keene, NH).

Samples for human mRNA isolation could include: peripheral blood mononuclear cells (monocytes, T cells, NK cells, granulocytes, B cells), resting (T100); peripheral blood mononuclear cells, activated with anti-CD3 for 2, 6, 12 h pooled (T101); T cell, TH0 clone Mot 72, resting (T102); T cell, TH0 clone Mot 72, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T103); T cell, TH0 clone Mot 72, anergic treated with specific peptide for 2, 7, 12 h pooled (T104); T cell, TH1 clone HY06, resting (T107); T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T108); T cell, TH1 clone HY06, anergic treated with specific peptide for 2, 6, 12 h pooled (T109); T cell, TH2 clone HY935, resting (T110); T cell, TH2 clone HY935, activated with anti-CD28 and anti-CD3 for 2, 7, 12 h pooled (T111); T cells CD4+CD45RO- T cells polarized 27 days in anti-CD28, IL-4, and anti IFN-γ, TH2 polarized, activated with anti-CD3 and anti-CD28 4 h (T116); T cell tumor lines Jurkat and Hut78, resting (T117); T cell clones, pooled AD130.2, Tc783.12, Tc783.13, Tc783.58, Tc782.69, resting (T118); T cell random γδ T cell clones, resting (T119); splenocytes, resting (B100); splenocytes, activated with anti-CD40 and IL-4 (B101); B cell EBV lines pooled WT49, RSB, JY, CVIR, 721.221, RM3, HSY, resting (B102); B cell line JY, activated with PMA and ionomycin for 1, 6 h pooled (B103); natural killer (NK) 20 clones pooled, resting (K100); NK 20 clones pooled, activated with PMA and ionomycin for 6 h (K101); NKL clone, derived from peripheral blood of LGL leukemia patient, IL-2 treated (K106); NK cytotoxic clone 640-A30-1, resting (K107); hematopoietic precursor line TF1, activated with PMA and ionomycin for 1, 6 h pooled (C100); U937 premonocytic line, resting (M100); U937 premonocytic line, activated with PMA and ionomycin for 1, 6 h pooled (M101); elutriated monocytes, activated with LPS, IFNγ,

anti-IL-10 for 1, 2, 6, 12, 24 h pooled (M102); elutriated monocytes, activated with LPS, IFN γ , IL-10 for 1, 2, 6, 12, 24 h pooled (M103); elutriated monocytes, activated with LPS, IFN γ , anti-IL-10 for 4, 16 h pooled (M106); elutriated

5 monocytes, activated with LPS, IFN γ , IL-10 for 4, 16 h pooled (M107); elutriated monocytes, activated LPS for 1 h (M108); elutriated monocytes, activated LPS for 6 h (M109); dendritic cells (DC) 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, resting (D101); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days,

10 activated with PMA and ionomycin for 1 hr (D102); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, activated with PMA and ionomycin for 6 hr (D103); DC 95% CD1a+, from CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D104); DC 95% CD14+, from CD34+ GM-CSF, TNF α 12

15 days FACS sorted, activated with PMA and ionomycin 1, 6 hr pooled (D105); DC CD1a+ CD86+, from CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D106); DC from monocytes GM-CSF, IL-4 5 days, resting (D107); DC from monocytes GM-CSF, IL-4 5 days, resting (D108);

20 DC from monocytes GM-CSF, IL-4 5 days, activated LPS 4, 16 h pooled (D109); DC from monocytes GM-CSF, IL-4 5 days, activated TNF α , monocyte supe for 4, 16 h pooled (D110); leiomyoma L11 benign tumor (X101); normal myometrium M5 (O115); malignant leiomyosarcoma GS1 (X103); lung fibroblast

25 sarcoma line MRC5, activated with PMA and ionomycin for 1, 6 h pooled (C101); kidney epithelial carcinoma cell line CHA, activated with PMA and ionomycin for 1, 6 h pooled (C102); kidney fetal 28 wk male (O100); lung fetal 28 wk male (O101); liver fetal 28 wk male (O102); heart fetal 28 wk male (O103);

30 brain fetal 28 wk male (O104); gallbladder fetal 28 wk male (O106); small intestine fetal 28 wk male (O107); adipose tissue fetal 28 wk male (O108); ovary fetal 25 wk female (O109); uterus fetal 25 wk female (O110); testes fetal 28 wk male (O111); spleen fetal 28 wk male (O112); adult placenta 28

35 wk (O113); and tonsil inflamed, from 12 year old (X100).

Quantitative PCR using IL-1 ξ specific primers on cDNA libraries showed low expression levels across libraries. No

significant signal above background was detected, except for
 low (+), medium (++), or high (+++) relative signals were
 detected across libraries: mononuclear cells, activated; T
 cell, TH0 clone Mot 72, resting; T cell, TH0 clone Mot 72,
 5 activated; T cell, TH0 clone Mot 72, anergic; T cell, TH1
 clone HY06, resting (++); T cell, TH1 clone HY06, activated
 (++); T cell, TH1 clone HY06, anergic; T cell, TH2 clone
 HY935, resting; T cell, TH2 clone HY935, activate"; T cells
 CD4+, TH2 polarized, activated; T cell lines Jurkat and Hut78,
 10 resting; T cell clones, pooled, resting; T cell $\gamma\delta$ clones,
 resting (+); CD28- T cell clone in pME (+); TR1 (Treg1) T cell
 clone (++); splenocytes, resting; splenocytes, activated; B
 cell EBV lines, resting; B cell line JY, activated; NK 20
 clones pooled, resting (+); NK 20 clones pooled, activated; NK
 15 cytotoxic clone, resting (+++); hematopoietic precursor line
 TF1, activated; U937 premonocytic line, resting; U937
 premonocytic line, activated; monocytes, LPS, IFN γ , anti-IL-10;
 monocytes, LPS, IFN γ , IL-10; monocytes, LPS, IFN γ , anti-IL-10,
 4 + 16 h (+); monocytes, LPS, IFN γ , IL-10, 4 + 16 h; monocytes,
 20 LPS, 1 h; monocytes, LPS, 6 h; DC 70% CD1a+, from CD34+ GM-
 CSF, TNF α , resting; DC 70% CD1a+, from CD34+ GM-CSF, TNF α ,
 activated 1 h; DC 70% CD1a+, from CD34+ GM-CSF, TNF α ,
 activated 6 h; DC 95% CD1a+, from CD34+ GM-CSF, TNF α ,
 activated 1 + 6 h; DC 95% CD14+, from CD34+ GM-CSF, TNF α ,
 25 activated 1 + 6 h; DC CD1a+ CD86+, from CD34+ GM-CSF, TNF α ,
 activated 1 + 6 h; DC ex monocytes GM-CSF, IL-4, resting; DC
 from monocytes GM-CSF, IL-4, resting; DC from monocytes GM-
 CSF, IL-4, LPS activated 4 + 16 h; DC from monocytes GM-CSF,
 IL-4, monokine activated 4 + 16 h; epithelial cells,
 30 unstimulated; epithelial cells, IL-1 β activated; lung
 fibroblast sarcoma line MRC5, activated; kidney epithelial
 carcinoma cell line CHA, activated; normal w.t. monkey lung;
 Ascaris-challenged monkey lung, 4 h; Ascaris-challenged monkey
 lung, 24 h; pool of two normal human lung samples; pool of
 35 three heavy smoker human lung samples; allergic lung sample;
 Pneumocystic carinii pneumonia lung sample; normal w.t. monkey
 colon; normal human colon; ulcerative colitis human colon

sample; normal human thyroid; Hashimoto's thyroiditis thyroid sample; pool of rheumatoid arthritis samples, human; normal human skin (+++); psoriasis patient skin sample (+++); tonsil inflamed; kidney fetal (++); lung fetal (+); liver fetal; heart fetal; brain fetal; gallbladder fetal (+); small intestine fetal (+); adipose tissue fetal; ovary fetal (+); uterus fetal; testes fetal; spleen fetal; placenta 28 wk (+); T cell, TH0 clone Mot 72, resting; T cell, TH0 clone Mot 72, activated; T cell, TH0 clone B21, resting; T cell, TH0 clone B21, activated; T cell, TH1 clone TA20-23, resting; T cell, TH1 clone TA20-23, activated; Jurkat cell line; and genomic DNA (++). Notably, expression was detected in normal skin, which is upregulated in psoriatic skin.

Because of the elevated signal in skin, various skin derived cell samples were evaluated by quantitative PCR. While all of the signals were, in absolute terms, quite low, no significant signal above background was detected, except for low (+) or medium (++) levels in: fibroblasts IL-10 18 h (+); fibroblasts IL-10 6 h; fibroblasts IL-4 18 h; fibroblasts IL-4 6 h; fibroblasts IFN γ 18 h; fibroblasts IFN γ 6 h; fibroblasts 3d pass untreated (+); keratinocytes IL-10 18 h (++); keratinocytes IL-10 6 h (++); keratinocytes IL-4 18 h (++); keratinocytes IL-4 6 h (++); keratinocytes IFN γ 18 h (++); keratinocytes IFN γ 6 h (++); keratinocytes 3d pass untreated (++); CLA $^{+}$ T cells; CLA $^{-}$ T cells; Langerhans cells (+); fibroblasts 7th passage (+); and keratinocytes 8th passage (+). These Langerhans and T cells were isolated from skin biopsies. In summary, the IL-1 ξ message is mainly expressed by the keratinocytes, and at lower levels in fibroblasts and Langerhans cells, with no significant expression in these T cells. The expression levels are comparable to that of the other IL-1 related messages IL-1 δ and IL-1 ϵ , each at some 10-100 fold lower than levels of expression of primate IL-1 α , IL-1 β , or IL-1 γ .

Likewise, in dendritic cells (DC), the signals were, in absolute terms, quite low. Dendritic cells (DC) are antigen-processing or presenting cells, and are found in all tissues

of the body. They can be classified into various categories, including: interstitial dendritic cells of the heart, kidney, gut, and lung; Langerhans cells in the skin and mucous membranes; interdigitating dendritic cells in the thymic medula and secondary lymphoid tissue; and blood and lymph dendritic cells. For a review of dendritic cells, see Steinman (1991) Annual Review of Immunology 9:271-296; and Banchereau and Schmitt (eds. 1994) Dendritic Cells in Fundamental and Clinical Immunology Plenum Press, NY.

Although dendritic cells in each of these compartments are CD45+ leukocytes that apparently arise from bone marrow, they may exhibit differences that relate to maturation state and microenvironment.

These dendritic cells again did not express large amounts of message. No significant signal above background was detected in pDC2 subset of dendritic cells (monocyte derived DC). Low message levels were detected in the DC1 (lymphocyte derived DC) day 0 and day 1 cultures. Higher levels were detected in the day 2, 3, 5, 6, 7, and 8 DC1 cultures.

Using the information described herein for cloning species variants, expression of rodent IL-1 ζ can be determined as above using, e.g., a murine homologue as for a detectable probe.

IV. Cloning of species counterparts of IL-1 ζ

Various strategies are used to obtain species counterparts of primate IL-1 ζ . One method is by cross hybridization using closely related species DNA probes. The degree of identity between mouse and human IL-1 counterparts typically is as high as 70%. It may be useful to go into evolutionarily similar species as intermediate steps. Another method is by using specific PCR primers based on the identification of blocks of similarity between human and mouse IL-1 counterparts, e.g., areas of highly conserved polypeptide sequence.

In addition, the IL-1 α , IL-1 β , and IL-1RA genes cluster on the same human chromosome. The fourth known members of the

IL-1 family, IL-1 γ , which is most closely related to IL-1 β , has been mapped to a different human chromosome. Duplication of the intact IL-1 α , IL-1 β , IL-1RA gene cluster, a potential genetic event explaining a proliferation of additional family members, would suggest the existence of two as yet unidentified IL-1 genes at the location of the IL-1 γ locus. IL-1 ζ is a potential candidate, and sequencing of the human IL-1 γ locus may well lead to identification of the novel IL-1 genes.

10

V. Production of mammalian IL-1 ζ protein

An appropriate, e.g., GST, fusion construct is engineered for expression, e.g., in *E. coli*. For example, a mouse IGIF pGex plasmid is constructed and transformed into *E. coli*. Freshly transformed cells are grown in LB medium containing 50 μ g/ml ampicillin and induced with IPTG (Sigma, St. Louis, MO). After overnight induction, the bacteria are harvested and the pellets containing IL-1 ζ are isolated. The pellets are homogenized in TE buffer (50 mM Tris-base pH 8.0, 10 mM EDTA and 2 mM pefabloc) in 2 liters. This material is passed through a microfluidizer (Microfluidics, Newton, MA) three times. The fluidized supernatant is spun down on a Sorvall GS-3 rotor for 1 h at 13,000 rpm. The resulting supernatant containing the IL-1 ζ is filtered and passed over a glutathione-SEPHAROSE column equilibrated in 50 mM Tris-base pH 8.0. The fractions containing the IL-1 ζ -GST fusion protein are pooled and cleaved with thrombin (Enzyme Research Laboratories, Inc., South Bend, IN). The cleaved pool is then passed over a Q-SEPHAROSE column equilibrated in 50 mM Tris-base. Fractions containing IL-1 ζ are pooled and diluted in cold distilled H₂O, to lower the conductivity, and passed back over a fresh Q-SEPHAROSE column. Fractions containing IL-1 ζ are pooled, aliquoted, and stored in the -70° C freezer.

Alternatively, the polypeptide may be produced in a non-fusion construct and expressed. The protein may be purified, either from secreted form, or from inclusion bodies, as appropriate.

35

Comparison of the CD spectrum with other IL-1 family members may suggest that the protein is correctly folded. See Hazuda, et al. (1969) J. Biol. Chem. 264:1689-1693.

Human IL-1 ζ epitope tagged protein gets secreted, e.g.,
5 from transiently transfected 293 T cells, as shown by presence of C-terminally tagged protein in the culture supernatant.

VI. Biological Assays with IL-1 ζ

Biological assays confirmed IFN- γ inducing activity by
10 IL-1 γ on T cells. IL-1 γ stimulates production of IFN- γ by purified NK cells, and that induction is strongly synergized with IL-12 or IL-2. Similar biological activity should be exhibited by IL-1 ζ or their antagonists.

The family of interleukins 1 contains molecules, each of
15 which is an important mediator of inflammatory disease. For a comprehensive review, see Dinarello (1996) "Biologic basis for interleukin-1 in disease" Blood 87:2095-2147. There are indications that the various IL-1's play important roles in the initiation of disease, including the recently identified
20 IGIF/IL-1 γ (e.g., Rothe, et al. (1997) "Active stage of autoimmune diabetes is associated with the expression of a novel cytokine, IGIF, which is located near Idd2." J. Clin. Invest. 99:469-474. The finding of novel proteins related to the IL-1 family furthers the identification of molecules that
25 provide the molecular basis for initiation of disease and allow for the development of therapeutic strategies of increased range and efficacy.

Similar biological assays as applied to other known members of the family should be performed with purified IL-
30 1 ζ .

VII. Preparation of antibodies specific for IL-1 ζ

Inbred Balb/c mice are immunized intraperitoneally with recombinant forms of the protein, e.g., purified soluble IL-
35 1 ζ -FLAG or stable transfected NIH-3T3 cells. Animals are boosted at appropriate time points with protein, with or without additional adjuvant, to further stimulate antibody

production. Serum is collected, or hybridomas produced with harvested spleens.

Alternatively, Balb/c mice are immunized with cells transformed with the gene or fragments thereof, either
5 endogenous or exogenous cells, or with isolated membranes enriched for expression of the antigen. Serum is collected at the appropriate time, typically after numerous further administrations. Various gene therapy techniques may be useful, e.g., in producing protein in situ, for generating an
10 immune response. Operable association of heterologous promoters with natural gene sequences is also provided, as are vectors encoding the DIRS3 with a receptor partner. See, e.g., Treco, et al. WO96/29411 or USSN 08/406,030.

Monoclonal antibodies may be made. For example,
15 splenocytes are fused with an appropriate fusion partner and hybridomas are selected in growth medium by standard procedures. Hybridoma supernatants are screened for the presence of antibodies which bind to the desired IL-1 ζ , e.g., by ELISA or other assay. Antibodies which specifically
20 recognize IL-1 ζ may also be selected or prepared, e.g., by immunoselection or related methods.

In another method, synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current
25 Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press. In appropriate situations, the binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods.
30 Nucleic acids may also be introduced into cells in an animal to produce the antigen, which serves to elicit an immune response. See, e.g., Wang, et al. (1993) Proc. Nat'l. Acad. Sci. 90:4156-4160; Barry, et al. (1994) BioTechniques 16:616-619; and Xiang, et al. (1995) Immunity 2: 129-135.

VIII. Production of fusion proteins with IL-1 ζ

Various fusion constructs are made with IL-1 ζ . This portion of the gene is fused to an epitope tag, e.g., a FLAG tag, or to a two hybrid system construct. See, e.g., Fields
5 and Song (1989) Nature 340:245-246.

The epitope tag may be used in an expression cloning procedure with detection with anti-FLAG antibodies to detect a binding partner, e.g., receptor for the IL-1 ζ . The two hybrid system may also be used to isolate proteins which specifically
10 bind to IL-1 ζ .

IX. Chromosome Mapping of IL-1 ζ

Chromosome spreads are prepared. In situ hybridization is performed on chromosome preparations obtained from
15 phytohemagglutinin-stimulated lymphocytes cultured for 72 h. 5-bromodeoxyuridine is added for the final seven hours of culture (60 μ g/ml of medium), to ensure a posthybridization chromosomal banding of good quality.

An appropriate fragment, e.g., a PCR fragment, is
20 amplified with the help of primers on total B cell cDNA template, and cloned into an appropriate vector. The vector is labeled by nick-translation with 3 H. The radiolabeled probe is hybridized to metaphase spreads, e.g., as described in Mattei, et al. (1985) Hum. Genet. 69:327-331.

25 After coating with nuclear track emulsion (KODAK NTB₂), slides are exposed, e.g., for 18 days at 4° C. To avoid slipping of silver grains during the banding procedure, chromosome spreads are first stained with buffered Giemsa solution and metaphase photographed. R-banding is then
30 performed by the fluorochrome-photolysis-Giemsa (FPG) method and metaphases rephotographed before analysis.

X. Structure activity relationship

Information on the criticality of particular residues is
35 determined using standard procedures and analysis. Standard mutagenesis analysis is performed, e.g., by generating many different variants at determined positions, e.g., at the

positions identified above, and evaluating biological activities of the variants. This may be performed to the extent of determining positions which modify activity, or to

e.g., of IL-1 γ -FLAG cDNA at 1 and 1/200 dilution, and a negative mock. Rinse cells with serum free DME. Add the DNA solution and incubate 5 hr at 37° C. Remove the medium and add 0.5 ml 10% DMSO in DME for 2.5 min. Remove and wash once
5 with DME. Add 1.5 ml growth medium and incubate overnight.

On day 2, change the medium. On days 3 or 4, the cells are fixed and stained. Rinse the cells twice with Hank's Buffered Saline Solution (HBSS) and fix in 4% paraformaldehyde (PFA)/glucose for 5 min. Wash 3X with HBSS. The slides may
10 be stored at -80° C after all liquid is removed. For each chamber, 0.5 ml incubations are performed as follows. Add HBSS/saponin (0.1%) with 32 μ l/ml of 1 M NaN₃ for 20 min. Cells are then washed with HBSS/saponin 1X. Add appropriate IL-1 δ or IL-1 δ /antibody complex to cells and incubate for 30
15 min. Wash cells twice with HBSS/saponin. If appropriate, add first antibody for 30 min. Add second antibody, e.g., Vector anti-mouse antibody, at 1/200 dilution, and incubate for 30 min. Prepare ELISA solution, e.g., Vector Elite ABC horseradish peroxidase solution, and preincubate for 30 min.
20 Use, e.g., 1 drop of solution A (avidin) and 1 drop solution B (biotin) per 2.5 ml HBSS/saponin. Wash cells twice with HBSS/saponin. Add ABC HRP solution and incubate for 30 min. Wash cells twice with HBSS, second wash for 2 min, which closes cells. Then add Vector diaminobenzoic acid (DAB) for 5
25 to 10 min. Use 2 drops of buffer plus 4 drops DAB plus 2 drops of H₂O₂ per 5 ml of glass distilled water. Carefully remove chamber and rinse slide in water. Air dry for a few minutes, then add 1 drop of Crystal Mount and a cover slip. Bake for 5 min at 85-90° C.

30 Evaluate positive staining of pools and progressively subclone to isolation of single genes responsible for the binding.

Alternatively, IL-1 ζ reagents are used to affinity purify or sort out cells expressing a receptor. See, e.g., Sambrook,
35 et al. or Ausubel, et al.

Another strategy is to screen for a membrane bound receptor by panning. The receptor cDNA is constructed as

described above. The ligand can be immobilized and used to immobilize expressing cells. Immobilization may be achieved by use of appropriate antibodies which recognize, e.g., a FLAG sequence of a IL-1 ζ fusion construct, or by use of antibodies
5 raised against the first antibodies. Recursive cycles of selection and amplification lead to enrichment of appropriate clones and eventual isolation of receptor expressing clones.

Phage expression libraries can be screened by mammalian IL-1 ζ . Appropriate label techniques, e.g., anti-FLAG
10 antibodies, will allow specific labeling of appropriate clones.

All citations herein are incorporated herein by reference to the same extent as if each individual publication or patent
15 application was specifically and individually indicated to be incorporated by reference including all figures and drawings.

Many modifications and variations of this invention, as will be apparent to one of ordinary skill in the art can be made to adapt to a particular situation, material, composition
20 of matter, process, process step or steps, to preserve the objective, spirit and scope of the invention. All such modifications are intended to be within the scope of the claims appended hereto without departing from the spirit and scope of the invention. The specific embodiments described
25 herein are offered by way of example only, and the invention is to be limited by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled; and the invention is not to be limited by the specific embodiments that have been presented herein by way of
30 example.

WHAT IS CLAIMED IS:

1. An isolated or recombinant polypeptide that:
 - a) specifically binds polyclonal antibodies generated
5 against at least a 12 consecutive amino acid segment
of SEQ ID NO: 2 or 4; and
 - b) comprises at least one sequence selected from:
 - i) GENSGVK; EDWEKD; CCLEDPA; FVHTSR; KKFSIHD;
VLVLDS; NLIAVP; FFALAS; SSASAEK; SLILLGV;
10 FCLYCDK; PSLQLK; KLMKLAAQ; FIFYRAQ; SRNMLES;
WFICTS; EPVGVT; or FSFQPVC (see SEQ ID NO: 2);
or
 - ii) FVHTSP; SPILLGV; or SWNMLES (see SEQ ID NO: 4).
- 15 2. The polypeptide of claim 1:
 - a) wherein said polypeptide comprises a plurality of
said sequences selected from said groups in section
b); or
 - b) which specifically binds to polyclonal antibodies
20 generated against an immunogen selected from SEQ ID
NO: 2 or 4.
 - c) wherein said 12 consecutive amino acid segment is
selected from: GVKMGSEDWEKD; AGSPLEPGPSLP;
SRKVKSLNPKKF; HDQDHKVLVLDS; NLIAVPDKNYIR;
25 FALASSLSSASA; GQSHPSLQLKKE; MKLAAQKESARR;
FYRAQVGSRNML; TSCNCNEPVGVT; FENRKHIEFSFQ; or
PVCKAEMSPSEV (see SEQ ID NO: 2); or AVSPLEPGPSLP;
SPKVKNLNPCKF; or FYRAQVGSWNML (see SEQ ID NO: 4).
- 30 3. The polypeptide of Claim 2, wherein said
polypeptide:
 - i) comprises a mature protein;
 - ii) lacks a post-translational modification;
 - iii) is from a primate, including a human;
 - 35 iv) is a natural allelic variant of IL-1 ζ ;
 - v) has a length at least about 30 amino acids;

- vi) exhibits at least two non-overlapping epitopes that are specific for a primate IL-1 ζ ;
- vii) exhibits a sequence identity over a length of at least about 20 amino acids to SEQ ID NO: 2 or 4;
- 5 viii) is not glycosylated;
- ix) has a molecular weight of at least 10 kD with natural glycosylation;
- x) is a synthetic polypeptide;
- xi) is attached to a solid substrate;
- 10 xii) is conjugated to another chemical moiety;
- xiii) is a 5-fold or less substitution from natural sequence; or
- xiv) is a deletion or insertion variant from a natural sequence.
- 15
4. A composition of matter comprising:
- a) a sterile polypeptide of Claim 2;
- b) said sterile polypeptide of Claim 2 and a carrier, wherein said carrier is:
- 20 i) an aqueous compound, including water, saline, and/or buffer; and/or
- ii) formulated for oral, rectal, nasal, topical, or parenteral administration.
- 25 5. A fusion protein having a polypeptide sequence of Claim 2 further comprising:
- a) a mature polypeptide of Claim 2;
- b) a detection or purification tag, including a FLAG, His6, or Ig sequence; or
- 30 c) sequence of another cytokine or chemokine.
6. A kit comprising a polypeptide of Claim 2, and:
- a) a compartment comprising said polypeptide; and/or
- b) instructions for use or disposal of reagents in said
- 35 kit.

7. A binding compound comprising an antigen binding site from an antibody, which specifically binds to a mature polypeptide of Claim 2, wherein:
- a) said mature polypeptide is a primate IL-1 ζ ;
 - 5 b) said binding compound is an Fv, Fab, or Fab2 fragment;
 - c) said binding compound is conjugated to another chemical moiety; or
 - d) said antibody:
 - 10 i) is raised against a 12 consecutive amino acid segment of SEQ ID NO: 2 or 4;
 - ii) is raised against a mature IL-1 ζ ;
 - iii) is raised to a purified primate IL-1 ζ ;
 - iv) is immunoselected;
 - 15 v) is a polyclonal antibody;
 - vi) binds to a denatured IL-1 ζ ;
 - vii) exhibits a Kd to antigen of at least 30 μ M;
 - viii) is attached to a solid substrate, including a bead or plastic membrane;
 - 20 ix) is in a sterile composition; or
 - x) is detectably labeled, including a radioactive or fluorescent label.
8. A kit comprising said binding compound of Claim 7,
- 25 and:
- a) a compartment comprising said binding compound; and/or
 - b) instructions for use or disposal of reagents in said kit.
- 30
9. A composition comprising:
- a) a sterile binding compound of Claim 7, or
 - b) said binding compound of Claim 7 and a carrier, wherein said carrier is:
 - 35 i) an aqueous compound, including water, saline, and/or buffer; and/or

- ii) formulated for oral, rectal, nasal, topical, or parenteral administration.

10. An isolated or recombinant nucleic acid encoding a polypeptide of Claim 2, wherein:
- a) said polypeptide of Claim 2 is a primate IL-1 ζ ; or
 - b) said nucleic acid:
 - i) encodes an antigenic peptide sequence of SEQ ID NO: 2 or 4;
 - 10 ii) encodes a plurality of antigenic peptide sequences of SEQ ID NO: 2 or 4;
 - iii) exhibits at least about 80% identity to a natural cDNA encoding said segment;
 - iv) is an expression vector;
 - 15 v) further comprises an origin of replication;
 - vi) is from a natural source;
 - vii) comprises a detectable label;
 - viii) comprises synthetic nucleotide sequence;
 - ix) is less than 6 kb, preferably less than 3 kb;
 - 20 x) is from a rodent;
 - xi) comprises a natural full length coding sequence;
 - xii) is a hybridization probe for a gene encoding said IL-1 ζ ; or
 - 25 xiii) is a PCR primer, PCR product, or mutagenesis primer; or
 - xiv) encodes an IL-1 ζ polypeptide.

11. A cell, transformed with said nucleic acid of Claim 10.

12. The cell of Claim 11, wherein said cell is:
- a) a prokaryotic cell;
 - b) a eukaryotic cell;
 - 35 c) a bacterial cell;
 - d) a yeast cell;
 - e) an insect cell;

- f) a mammalian cell;
 - g) a mouse cell;
 - h) a primate cell; or
 - i) a human cell.
-

18. The method of Claim 17, wherein:
- a) said contacting is in combination with an agonist or antagonist of IL-1 α , IL-1RA, IL-1 β , IL-1 γ , IL-1 δ , IL-1 ϵ , IL-2, and/or IL-12;
 - 5 b) said contacting is with an antagonist, including binding composition comprising an antibody binding site which specifically binds an IL-1 ζ ; or
 - c) said modulating is regulation of IFN- γ production.
- 10 19. A binding compound comprising an antigen binding portion from an antibody, which specifically binds to a primate protein of Claim 1, wherein:
- a) said protein is a human protein;
 - b) said binding compound is an Fv, Fab, or Fab2
 - 15 fragment;
 - c) said binding compound is conjugated to another chemical moiety; or
 - d) said antibody:
 - i) is raised against a polypeptide sequence of a
 - 20 mature polypeptide comprising at least 12 consecutive amino acids of SEQ ID NO: 2 or 4;
 - ii) is raised against a mature primate IL-1 ζ ;
 - iii) is raised to a purified primate IL-1 ζ ;
 - iv) is immunoselected;
 - 25 v) is a polyclonal antibody;
 - vi) binds to a denatured primate IL-1 ζ ;
 - vii) exhibits a Kd to antigen of at least 30 μ M;
 - viii) is attached to a solid substrate, including a bead or plastic membrane;
 - 30 ix) is in a sterile composition; or
 - x) is detectably labeled, including a radioactive or fluorescent label.

20. A method of:
- A) making an antibody of Claim 19, comprising immunizing an immune system with an immunogenic amount of:
 - a) a primate IL-1 ζ polypeptide; or
 - 5 b) a peptide sequence comprising at least 12 consecutive amino acids of SEQ ID NO: 2 or 4; thereby causing said antibody to be produced; or
 - B) producing an antigen:antibody complex, comprising contacting a primate IL-1 ζ polypeptide with an
10 antibody of Claim 19 thereby allowing said complex to form.

SEQUENCE SUBMISSION

SEQ ID NO: 1 provides primate IL-1 ζ nucleotide sequence.
 SEQ ID NO: 2 provides primate IL-1 ζ polypeptide sequence.
 5 SEQ ID NO: 3 provides primate IL-1 ζ variant nucleotide sequence.
 SEQ ID NO: 4 provides primate IL-1 ζ variant polypeptide sequence.
 SEQ ID NO: 5 provides primate IL-1 α polypeptide sequence.
 SEQ ID NO: 6 provides rodent IL-1 α polypeptide sequence.
 10 SEQ ID NO: 7 provides primate IL-1 γ polypeptide sequence.
 SEQ ID NO: 8 provides rodent IL-1 γ polypeptide sequence.
 SEQ ID NO: 9 provides primate IL-1 β polypeptide sequence.
 SEQ ID NO: 10 provides rodent IL-1 β polypeptide sequence.
 SEQ ID NO: 11 provides primate IL-1RA polypeptide sequence.
 SEQ ID NO: 12 provides rodent IL-1RA polypeptide sequence.
 15 SEQ ID NO: 13 provides rodent IL-1 δ polypeptide sequence.
 SEQ ID NO: 14 provides rodent IL-1 ϵ polypeptide sequence.
 SEQ ID NO: 15 provides primate IL-1 ϵ polypeptide sequence.

20 <110> Timans, Jacqueline C.
 <120> Mammalian Cytokines; Related Reagents and Methods
 <130> DX0904K

25
 <140>
 <141>

30 <150> 09/156,966
 <151> 1998-09-18
 <160> 15
 <170> PatentIn Ver. 2.1

35 <210> 1
 <211> 1225
 <212> DNA
 <213> primate

40
 <220>
 <221> CDS
 <222> (491)..(1144)

45 <400> 1
 CGGTTTGTTT TCTTTAGAGA TTTTACAGTG TTGGTTATAA TTGTGCACTT AATCTTTATT 60
 TTCCTTATAC AGTAGTCCCC CCGATCAACT GGGGGCATGT TCCATACCCC TGGTGGATTC 120
 50 CTGAAACTGC CAGTTAGTAC CAAACCCTAT ATAGATTGTG TTTTTCCTG TACGCAGGCC 180
 GACACACAGG AAATCATAAG TCAGGAGGGC CACTGCCACG CAGGAAAGAC CCATCTGAAC 240
 TGCTGCAAAA GCTCCGTGTC GATTTATTGC TTCCACAAAT AGTGCCGATA TGCACCAGGC 300
 55 ACTGTTGTAA AACTGAAAAT ATGTTTTGGG ATGTGCCAG TCTACCTAGC TCCTTCAAGT 360
 AAAGGATCCT GAGAACTGAA GGCAAACAGA GCTCCAGGAG TCCAAGACAG AGCCACACAC 420
 60 CACGAGGATC CTGGCCCAGG TCTTGGACTT CCATTCCCAT TTCTGTTGAG TAATAAACTC 480
 AACGTTGAAA ATG TCC TTT GTG GGG GAG AAC TCA GGA GTG AAA ATG GGC 529

	Met	Ser	Phe	Val	Gly	Glu	Asn	Ser	Gly	Val	Lys	Met	Gly				
	1				5					10							
5	TCT	GAG	GAC	TGG	GAA	AAA	GAT	GAA	CCC	CAG	TGC	TGC	TTA	GAA	GAC	CCG	577
	Ser	Glu	Asp	Trp	Glu	Lys	Asp	Glu	Pro	Gln	Cys	Cys	Leu	Glu	Asp	Pro	
	15						20					25					
10	GCT	GGA	AGC	CCC	CTG	GAA	CCA	GGC	CCA	AGC	CTC	CCC	ACC	ATG	AAT	TTT	625
	Ala	Gly	Ser	Pro	Leu	Glu	Pro	Gly	Pro	Ser	Leu	Pro	Thr	Met	Asn	Phe	
	30					35					40					45	
15	GTT	CAC	ACA	AGT	CGA	AAG	GTG	AAG	AGC	TTA	AAC	CCG	AAG	AAA	TTC	AGC	673
	Val	His	Thr	Ser	Arg	Lys	Val	Lys	Ser	Leu	Asn	Pro	Lys	Lys	Phe	Ser	
					50					55					60		
20	ATT	CAT	GAC	CAG	GAT	CAC	AAA	GTA	CTG	GTC	CTG	GAC	TCT	GGG	AAT	CTC	721
	Ile	His	Asp	Gln	Asp	His	Lys	Val	Leu	Val	Leu	Asp	Ser	Gly	Asn	Leu	
				65				70						75			
25	ATA	GCA	GTT	CCA	GAT	AAA	AAC	TAC	ATA	CGC	CCA	GAG	ATC	TTC	TTT	GCA	769
	Ile	Ala	Val	Pro	Asp	Lys	Asn	Tyr	Ile	Arg	Pro	Glu	Ile	Phe	Phe	Ala	
			80					85					90				
30	TTA	GCC	TCA	TCC	TTG	AGC	TCA	GCC	TCT	GCG	GAG	AAA	GGA	AGT	CTG	ATT	817
	Leu	Ala	Ser	Ser	Leu	Ser	Ser	Ala	Ser	Ala	Glu	Lys	Gly	Ser	Leu	Ile	
		95					100					105					
35	CTC	CTG	GGG	GTC	TCT	AAA	GGG	GAG	TTT	TGT	CTC	TAC	TGT	GAC	AAG	GAT	865
	Leu	Leu	Gly	Val	Ser	Lys	Gly	Glu	Phe	Cys	Leu	Tyr	Cys	Asp	Lys	Asp	
	110					115					120					125	
40	AAA	GGA	CAA	AGT	CAT	CCA	TCC	CTT	CAG	CTG	AAG	AAG	GAG	AAA	CTG	ATG	913
	Lys	Gly	Gln	Ser	His	Pro	Ser	Leu	Gln	Leu	Lys	Lys	Glu	Lys	Leu	Met	
					130					135					140		
45	AAG	CTG	GCT	GCC	CAA	AAG	GAA	TCA	GCA	CGC	CGG	CCC	TTC	ATC	TTT	TAT	961
	Lys	Leu	Ala	Ala	Gln	Lys	Glu	Ser	Ala	Arg	Arg	Pro	Phe	Ile	Phe	Tyr	
				145				150					155				
50	AGG	GCT	CAG	GTG	GGC	TCC	CGG	AAC	ATG	CTG	GAG	TCG	GCG	GCT	CAC	CCC	1009
	Arg	Ala	Gln	Val	Gly	Ser	Arg	Asn	Met	Leu	Glu	Ser	Ala	Ala	His	Pro	
			160					165					170				
55	GGA	TGG	TTC	ATC	TGC	ACC											

3

	Met	Ser	Phe	Val	Gly	Glu	Asn	Ser	Gly	Val	Lys	Met	Gly	Ser	Glu	Asp	
	1				5					10					15		
5	Trp	Glu	Lys	Asp	Glu	Pro	Gln	Cys	Cys	Leu	Glu	Asp	Pro	Ala	Gly	Ser	
				20					25					30			
	Pro	Leu	Glu	Pro	Gly	Pro	Ser	Leu	Pro	Thr	Met	Asn	Phe	Val	His	Thr	
			35					40					45				
10	Ser	Arg	Lys	Val	Lys	Ser	Leu	Asn	Pro	Lys	Lys	Phe	Ser	Ile	His	Asp	
		50					55					60					
	Gln	Asp	His	Lys	Val	Leu	Val	Leu	Asp	Ser	Gly	Asn	Leu	Ile	Ala	Val	
15		65				70				75						80	
	Pro	Asp	Lys	Asn	Tyr	Ile	Arg	Pro	Glu	Ile	Phe	Phe	Ala	Leu	Ala	Ser	
				85						90					95		
20	Ser	Leu	Ser	Ser	Ala	Ser	Ala	Glu	Lys	Gly	Ser	Leu	Ile	Leu	Leu	Gly	
				100					105					110			
	Val	Ser	Lys	Gly	Glu	Phe	Cys	Leu	Tyr	Cys	Asp	Lys	Asp	Lys	Gly	Gln	
			115					120					125				
25	Ser	His	Pro	Ser	Leu	Gln	Leu	Lys	Lys	Glu	Lys	Leu	Met	Lys	Leu	Ala	
		130					135					140					
	Ala	Gln	Lys	Glu	Ser	Ala	Arg	Arg	Pro	Phe	Ile	Phe	Tyr	Arg	Ala	Gln	
30		145				150					155					160	
	Val	Gly	Ser	Arg	Asn	Met	Leu	Glu	Ser	Ala	Ala	His	Pro	Gly	Trp	Phe	
				165						170					175		
35	Ile	Cys	Thr	Ser	Cys	Asn	Cys	Asn	Glu	Pro	Val	Gly	Val	Thr	Asp	Lys	
				180					185					190			
	Phe	Glu	Asn	Arg	Lys	His	Ile	Glu	Phe	Ser	Phe	Gln	Pro	Val	Cys	Lys	
			195					200					205				
40	Ala	Glu	Met	Ser	Pro	Ser	Glu	Val	Ser	Asp							
		210					215										
45	<210> 3																
	<211> 657																
	<212> DNA																
	<213> primate																
50	<220>																
	<221> CDS																
	<222> (1) .. (654)																
55	<400> 3																
	ATG	TCC	TTT	GTG	GGG	GAG	AAC	TCA	GGA	GTG	AAA	ATG	GGC	TCT	GAG	GAC	48
	Met	Ser	Phe	Val	Gly	Glu	Asn	Ser	Gly	Val	Lys	Met	Gly	Ser	Glu	Asp	
	1				5					10					15		
60	TGG	GAA	AAA	GAT	GAA	CCC	CAG	TGC	TGC	TTA	GAA	GAC	CCG	GCT	GTA	AGC	96
	Trp	Glu	Lys	Asp	Glu	Pro	Gln	Cys	Cys	Leu	Glu	Asp	Pro	Ala	Val	Ser	
				20					25					30			
65	CCC	CTG	GAA	CCA	GGC	CCA	AGC	CTC	CCC	GCC	ATG	AAT	TTT	GTT	CAC	ACA	144
	Pro	Leu	Glu	Pro	Gly	Pro	Ser	Leu	Pro	Ala	Met	Asn	Phe	Val	His	Thr	
			35					40					45				
	AGT	CCA	AAG	GTG	AAG	AAC	TTA	AAC	CCG	AAG	AAA	TTC	AGC	ATT	CAT	GAC	192

	Ser	Pro	Lys	Val	Lys	Asn	Leu	Asn	Pro	Lys	Lys	Phe	Ser	Ile	His	Asp	
	50						55					60					
5	CAG	GAT	CAC	AAA	GTA	CTG	GTC	CTG	GAC	TCT	GGG	AAT	CTC	ATA	GCA	GTT	240
	Gln	Asp	His	Lys	Val	Leu	Val	Leu	Asp	Ser	Gly	Asn	Leu	Ile	Ala	Val	
	65					70					75					80	
10	CCA	GAT	AAA	AAC	TAC	ATA	CGC	CCA	GAG	ATC	TTC	TTT	GCA	TTA	GCC	TCA	288
	Pro	Asp	Lys	Asn	Tyr	Ile	Arg	Pro	Glu	Ile	Phe	Phe	Ala	Leu	Ala	Ser	
					85					90					95		
15	TCC	TTG	AGC	TCA	GCC	TCT	GCG	GAG	AAA	GGA	AGT	CCG	ATT	CTC	CTG	GGG	336
	Ser	Leu	Ser	Ser	Ala	Ser	Ala	Glu	Lys	Gly	Ser	Pro	Ile	Leu	Leu	Gly	
				100					105					110			
20	GTC	TCT	AAA	GGG	GAG	TTT	TGT	CTC	TAC	TGT	GAC	AAG	GAT	AAA	GGA	CAA	384
	Val	Ser	Lys	Gly	Glu	Phe	Cys	Leu	Tyr	Cys	Asp	Lys	Asp	Lys	Gly	Gln	
			115				120					125					
25	AGT	CAT	CCA	TCC	CTT	CAG	CTG	AAG	AAG	GAG	AAA	CTG	ATG	AAG	CTG	GCT	432
	Ser	His	Pro	Ser	Leu	Gln	Leu	Lys	Lys	Glu	Lys	Leu	Met	Lys	Leu	Ala	
		130					135					140					
30	GCC	CAA	AAG	GAA	TCA	GCA	CGC	CGG	CCC	TTC	ATC	TTT	TAT	AGG	GCT	CAG	480
	Ala	Gln	Lys	Glu	Ser	Ala	Arg	Arg	Pro	Phe	Ile	Phe	Tyr	Arg	Ala	Gln	
	145					150					155					160	
35	GTG	GGC	TCC	TGG	AAC	ATG	CTG	GAG	TCG	GCG	GCT	CAC	CCC	GGA	TGG	TTC	528
	Val	Gly	Ser	Trp	Asn	Met	Leu	Glu	Ser	Ala	Ala	His	Pro	Gly	Trp	Phe	
					165					170					175		
40	ATC	TGC	ACC	TCC	TGC	AAT	TGT	AAT	GAG	CCT	GTT	GGG	GTG	ACA	GAT	AAA	576
	Ile	Cys	Thr	Ser	Cys	Asn	Cys	Asn	Glu	Pro	Val	Gly	Val	Thr	Asp	Lys	
				180					185				190				
45	TTT	GAG	AAC	AGG	AAA	CAC	ATT	GAA	TTT	TCA	TTT	CAA	CCA	GTT	TGC	AAA	624
	Phe	Glu	Asn	Arg	Lys	His	Ile	Glu	Phe	Ser	Phe	Gln	Pro	Val	Cys	Lys	
			195				200					205					
50	GCT	GAA	ATG	AGC	CCC	AGT	GAG	GTC	AGC	GAT	TAG						657
	Ala	Glu	Met	Ser	Pro	Ser	Glu	Val	Ser	Asp							
		210					215										
45	<210> 4																
	<211> 218																
	<212> PRT																
	<213> primate																
50	<400> 4																
	Met	Ser	Phe	Val	Gly	Glu	Asn	Ser	Gly	Val	Lys	Met	Gly	Ser	Glu	Asp	
	1				5					10					15		
55	Trp	Glu	Lys	Asp	Glu	Pro	Gln	Cys	Cys	Leu	Glu	Asp	Pro	Ala	Val	Ser	
				20					25					30			
	Pro	Leu	Glu	Pro	Gly	Pro	Ser	Leu	Pro	Ala	Met	Asn	Phe	Val	His	Thr	
			35					40					45				
60	Ser	Pro	Lys	Val	Lys	Asn	Leu	Asn	Pro	Lys	Lys	Phe	Ser	Ile	His	Asp	
		50					55					60					
65	Gln	Asp	His	Lys	Val	Leu	Val	Leu	Asp	Ser	Gly	Asn	Leu	Ile	Ala	Val	
	65					70					75					80	
	Pro	Asp	Lys	Asn	Tyr	Ile	Arg	Pro	Glu	Ile	Phe	Phe	Ala	Leu	Ala	Ser	

5

	85	90	95
5	Ser Leu Ser Ser Ala Ser Ala Glu Lys Gly Ser Pro Ile Leu Leu Gly		
	100	105	110
	Val Ser Lys Gly Glu Phe Cys Leu Tyr Cys Asp Lys Asp Lys Gly Gln		
	115	120	125
10	Ser His Pro Ser Leu Gln Leu Lys Lys Glu Lys Leu Met Lys Leu Ala		
	130	135	140
	Ala Gln Lys Glu Ser Ala Arg Arg Pro Phe Ile Phe Tyr Arg Ala Gln		
	145	150	155
15	Val Gly Ser Trp Asn Met Leu Glu Ser Ala Ala His Pro Gly Trp Phe		
	165	170	175
	Ile Cys Thr Ser Cys Asn Cys Asn Glu Pro Val Gly Val Thr Asp Lys		
	180	185	190
20	Phe Glu Asn Arg Lys His Ile Glu Phe Ser Phe Gln Pro Val Cys Lys		
	195	200	205
25	Ala Glu Met Ser Pro Ser Glu Val Ser Asp		
	210	215	
30	<210> 5		
	<211> 159		
	<212> PRT		
	<213> primate		
35	<400> 5		
	Ser Ala Pro Phe Ser Phe Leu Ser Asn Val Lys Tyr Asn Phe Met Arg		
	1	5	10
	Ile Ile Lys Tyr Glu Phe Ile Leu Asn Asp Ala Leu Asn Gln Ser Ile		
	20	25	30
40	Ile Arg Ala Asn Asp Gln Tyr Leu Thr Ala Ala Ala Leu His Asn Leu		
	35	40	45
45	Asp Glu Ala Val Lys Phe Asp Met Gly Ala Tyr Lys Ser Ser Lys Asp		
	50	55	60
	Asp Ala Lys Ile Thr Val Ile Leu Arg Ile Ser Lys Thr Gln Leu Tyr		
	65	70	75
50	Val Thr Ala Gln Asp Glu Asp Gln Pro Val Leu Leu Lys Glu Met Pro		
	85	90	95
	Glu Ile Pro Lys Thr Ile Thr Gly Ser Glu Thr Asn Leu Leu Phe Phe		
	100	105	110
55	Trp Glu Thr His Gly Thr Lys Asn Tyr Phe Thr Ser Val Ala His Pro		
	115	120	125
60	Asn Leu Phe Ile Ala Thr Lys Gln Asp Tyr Trp Val Cys Leu Ala Gly		
	130	135	140
	Gly Pro Pro Ser Ile Thr Asp Phe Gln Ile Leu Glu Asn Gln Ala		
	145	150	155
65	<210> 6		
	<211> 156		

6

<212> PRT
<213> rodent

<400> 6
5 Ser Ala Pro Tyr Thr Tyr Gln Ser Asp Leu Arg Tyr Lys Leu Met Lys
1 5 10 15
Leu Val Arg Gln Lys Phe Val Met Asn Asp Ser Leu Asn Gln Thr Ile
20 25 30
10 Tyr Gln Asp Val Asp Lys His Tyr Leu Ser Thr Thr Trp Leu Asn Asp
35 40 45
Leu Gln Gln Glu Val Lys Phe Asp Met Tyr Ala Tyr Ser Ser Gly Gly
15 50 55 60
Asp Asp Ser Lys Tyr Pro Val Thr Leu Lys Ile Ser Asp Ser Gln Leu
65 70 75 80
20 Phe Val Ser Ala Gln Gly Glu Asp Gln Pro Val Leu Leu Lys Glu Leu
85 90 95
Pro Glu Thr Pro Lys Leu Ile Thr Gly Ser Glu Thr Asp Leu Ile Phe
100 105 110
25 Phe Trp Lys Ser Ile Asn Ser Lys Asn Tyr Phe Thr Ser Ala Ala Tyr
115 120 125
Pro Glu Leu Phe Ile Ala Thr Lys Glu Gln Ser Arg Val His Leu Ala
130 135 140
30 Arg Gly Leu Pro Ser Met Thr Asp Phe Gln Ile Ser
145 150 155
35
<210> 7
<211> 158
<212> PRT
<213> primate
40
<400> 7
Asp Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu
1 5 10 15
45 Asn Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu
20 25 30
Asp Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe
35 40 45
50 Ile Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr
50 55 60
Ile Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys
55 65 70 75 80
Ile Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr
85 90 95
60 Lys Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn
100 105 110
Lys Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys
115 120 125
65 Glu Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu

	130		135		140
5	Leu Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp				
	145		150		155
	<210> 8				
	<211> 158				
	<212> PRT				
10	<213> rodent				
	<400> 8				
15	Asp Asn Phe Gly Arg Leu His Cys Thr Thr Ala Val Ile Arg Asn Ile				
	1		5		10
					15
	Asn Asp Gln Val Leu Phe Val Asp Lys Arg Gln Pro Val Phe Glu Asp				
		20		25	30
20	Met Thr Asp Ile Asp Gln Ser Ala Ser Glu Pro Gln Thr Arg Leu Ile				
		35		40	45
	Ile Tyr Met Tyr Lys Asp Ser Glu Val Arg Gly Leu Ala Val Thr Leu				
		50		55	60
25	Ser Val Lys Asp Ser Lys Met Ser Thr Leu Ser Cys Lys Asn Lys Ile				
		65		70	75
					80
	Ile Ser Phe Glu Glu Met Asp Pro Pro Glu Asn Ile Asp Asp Ile Gln				
		85		90	95
30	Ser Asp Leu Ile Phe Phe Gln Lys Arg Val Pro Gly His Asn Lys Met				
		100		105	110
	Glu Phe Glu Ser Ser Leu Tyr Glu Gly His Phe Leu Ala Cys Gln Lys				
35		115		120	125
	Glu Asp Asp Ala Phe Lys Leu Ile Leu Lys Lys Lys Asp Glu Asn Gly				
		130		135	140
40	Asp Lys Ser Val Met Phe Thr Leu Thr Asn Leu His Gln Ser				
		145		150	155
	<210> 9				
45	<211> 154				
	<212> PRT				
	<213> primate				
	<400> 9				
50	Asp Ala Pro Val Arg Ser Leu Asn Cys Thr Leu Arg Asp Ser Gln Gln				
		1		5	10
					15
	Lys Ser Leu Val Met Ser Gly Pro Tyr Glu Leu Lys Ala Leu His Leu				
		20		25	30
55	Gln Gly Gln Asp Met Glu Gln Gln Val Val Phe Ser Met Ser Phe Val				
		35		40	45
	Gln Gly Glu Glu Ser Asn Asp Lys Ile Pro Val Ala Leu Gly Leu Lys				
60		50		55	60
	Glu Lys Asn Leu Tyr Leu Ser Cys Val Leu Lys Asp Asp Lys Pro Thr				
		65		70	75
					80
65	Leu Gln Leu Glu Ser Val Asp Pro Lys Asn Tyr Pro Lys Lys Lys Met				
		85		90	95

Glu Lys Arg Phe Val Phe Asn Lys Ile Glu Ile Asn Asn Lys Leu Glu
 100 105 110
 5 Phe Glu Ser Ala Gln Phe Pro Asn Trp Tyr Ile Ser Thr Ser Gln Ala
 115 120 125
 Glu Asn Met Pro Val Phe Leu Gly Gly Thr Lys Gly Gly Gln Asp Ile
 130 135 140
 10 Thr Asp Phe Thr Met Gln Phe Val Ser Ser
 145 150
 15 <210> 10
 <211> 153
 <212> PRT
 <213> rodent
 20 <400> 10
 Asp Val Pro Ile Arg Gln Leu His Tyr Arg Leu Arg Asp Glu Gln Gln
 1 5 10 15
 25 Lys Ser Leu Val Leu Ser Asp Pro Tyr Glu Leu Lys Ala Leu His Leu
 20 25 30
 Asn Gly Gln Asn Ile Asn Gln Gln Val Ile Phe Ser Met Ser Phe Val
 35 40 45
 30 Gln Gly Glu Pro Ser Asn Asp Lys Ile Pro Val Ala Leu Gly Leu Lys
 50 55 60
 Gly Lys Asn Leu Tyr Leu Ser Cys Val Met Lys Asp Gly Thr Pro Thr
 65 70 75 80
 35 Leu Gln Leu Glu Ser Val Asp Pro Lys Gln Tyr Pro Lys Lys Lys Met
 85 90 95
 40 Glu Lys Arg Phe Val Phe Asn Lys Ile Glu Val Lys Ser Lys Val Glu
 100 105 110
 Phe Glu Ser Ala Glu Phe Pro Asn Trp Tyr Ile Ser Thr Ser Gln Ala
 115 120 125
 45 Glu His Lys Pro Val Phe Leu Gly Asn Asn Ser Gly Gln Asp Ile Ile
 130 135 140
 Asp Phe Thr Met Glu Ser Val Ser Ser
 145 150
 50
 <210> 11
 <211> 153
 <212> PRT
 55 <213> primate
 <400> 11
 Cys Arg Pro Ser Gly Arg Lys Ser Ser Lys Met Gln Ala Phe Arg Ile
 1 5 10 15
 60 Trp Asp Val Asn Gln Lys Thr Phe Tyr Leu Arg Asn Asn Gln Leu Val
 20 25 30
 65 Ala Gly Tyr Leu Gln Gly Pro Asn Val Asn Leu Glu Glu Lys Ile Asp
 35 40 45

9

Val Val Pro Ile Glu Pro His Ala Leu Phe Leu Gly Ile His Gly Gly
 50 55 60
 5 Lys Leu Cys Leu Ser Cys Val Lys Ser Gly Asp Glu Thr Arg Leu Gln
 65 70 75 80
 Leu Glu Ala Val Asn Ile Thr Asp Leu Ser Glu Asn Arg Lys Gln Asp
 85 90 95
 10 Lys Arg Phe Ala Phe Ile Arg Ser Asp Ser Gly Pro Thr Thr Ser Phe
 100 105 110
 Glu Ser Ala Ala Cys Pro Gly Trp Phe Leu Cys Thr Ala Met Glu Ala
 115 120 125
 15 Asp Gln Pro Val Ser Leu Thr Asn Met Pro Asp Glu Gly Val Met Val
 130 135 140
 20 Thr Lys Phe Tyr Phe Gln Glu Asp Glu
 145 150
 <210> 12
 <211> 153
 25 <212> PRT
 <213> rodent
 <400> 12
 30 Cys Arg Pro Ser Gly Lys Arg Pro Cys Lys Met Gln Ala Phe Arg Ile
 1 5 10 15
 Trp Asp Thr Asn Gln Lys Thr Phe Tyr Leu Arg Asn Asn Gln Leu Ile
 20 25 30
 35 Ala Gly Tyr Leu Gln Gly Pro Asn Ile Lys Leu Glu Glu Lys Ile Asp
 35 40 45
 Met Val Pro Ile Asp Leu His Ser Val Phe Leu Gly Ile Lys Gly Tyr
 50 55 60
 40 Lys Leu Tyr Met Ser Cys Val Lys Ser Gly Asp Asp Ile Lys Leu Gln
 65 70 75 80
 Leu Glu Glu Val Asn Ile Thr Asp Leu Ser Lys Asn Lys Glu Glu Asp
 85 90 95
 Lys Arg Phe Thr Phe Ile Arg Ser Glu Lys Gly Pro Thr Thr Ser Phe
 100 105 110
 50 Glu Ser Ala Ala Cys Pro Gly Trp Phe Leu Cys Thr Thr Leu Glu Ala
 115 120 125
 Asp Arg Pro Val Ser Leu Thr Asn Thr Pro Glu Glu Pro Leu Ile Val
 130 135 140
 55 Thr Lys Phe Tyr Phe Gln Glu Asp Gln
 145 150
 60 <210> 13
 <211> 156
 <212> PRT
 <213> rodent
 65 <400> 13
 Met Met Val Leu Ser Gly Ala Leu Cys Phe Arg Met Lys Asp Ser Ala

10

	1		5		10		15
	Leu	Lys	Val	Leu	Tyr	Leu	His
			20			25	
	Asn	Asn	Gln	Leu	Leu	Ala	Gly
			30				
5	His	Ala	Glu	Lys	Val	Ile	Lys
		35				40	
	Gly	Glu	Glu	Ile	Ser	Val	Val
					45		
	Pro	Asn					
10	Arg	Ala	Leu	Asp	Ala	Ser	Leu
		50				55	
	Ser	Pro	Val	Ile	Leu	Gly	Val
					60		
	Gln	Gly					
	Gly	Ser	Gln	Cys	Leu	Ser	Cys
		65			70		
	Gly	Thr	Glu	Lys	Gly	Pro	Ile
				75			
	Leu	Lys					
							80
15	Leu	Glu	Pro	Val	Asn	Ile	Met
				85			
	Glu	Leu	Tyr	Leu	Gly	Ala	Lys
			90				
	Glu	Ser					
	Lys	Ser	Phe	Thr	Phe	Tyr	Arg
			100				
	Arg	Arg	Asp	Met	Gly	Leu	Thr
			105				
20	Ser	Ser	Phe				
	Glu	Ser	Ala	Ala	Tyr	Pro	Gly
			115				
	Trp	Phe	Leu	Cys	Thr	Ser	Pro
	Glu	Ala					
	Asp	Gln	Pro	Val	Arg	Leu	Thr
		130				135	
25	Gln	Ile	Pro	Glu	Asp	Pro	Ala
					140		
	Trp	Asp					
	Ala	Pro	Ile	Thr	Asp	Phe	Tyr
					150		
	Gln	Gln	Cys	Asp			
			155				
30							
	<210>	14					
	<211>	160					
	<212>	PRT					
	<213>	rodent					
35							
	<400>	14					
	Met	Asn	Lys	Glu	Lys	Glu	Leu
		1			5		
	Arg	Ala	Ala	Ser	Pro	Ser	Leu
			10				
	Arg	His					
		15					
40	Val	Gln	Asp	Leu	Ser	Ser	Arg
			20				
	Val	Trp	Ile	Leu	Gln	Asn	Asn
		25				30	
	Ile	Leu					
	Thr	Ala	Val	Pro	Arg	Lys	Glu
			35				
	Gln	Thr	Val	Pro	Val	Thr	Ile
						45	
45	Leu	Pro	Cys	Gln	Tyr	Leu	Asp
		50				55	
	Thr	Leu	Glu	Thr	Asn	Arg	Gly
					60		
	Asp	Pro					
	Thr	Tyr	Met	Gly	Val	Gln	Arg
		65			70		
	Pro	Met	Ser	Cys	Leu	Phe	Cys
				75			
50	Thr	Lys					
							80
	Asp	Gly	Glu	Gln	Pro	Val	Leu
				85			
	Gln	Leu	Gly	Glu	Gly	Asn	Ile
			90				
	Met	Glu					
	Met	Tyr	Asn	Lys	Lys	Glu	Pro
			100				
	Val	Lys	Ala	Ser	Leu	Phe	Tyr
			105				
	His	Lys					
	Lys	Ser	Gly	Thr	Thr	Ser	Thr
			115				
	Phe	Glu	Ser	Ala	Ala	Phe	Pro
						125	
60	Gly	Trp					
	Phe	Ile	Ala	Val	Cys	Ser	Lys
						135	
	Gly	Ser	Cys	Pro	Leu	Ile	Leu
					140		
	Thr	Gln					
	Glu	Leu	Gly	Glu	Ile	Phe	Ile
		145				150	
65	Thr	Asp	Phe	Glu	Met	Ile	Val
				155			
	Val	Val	His				
			160				

11

<210> 15
 <211> 169
 <212> PRT
 5 <213> primate

 <400> 15
 Met Arg Gly Thr Pro Gly Asp Ala Asp Gly Gly Gly Arg Ala Val Tyr
 1 5 10 15
 10 Gln Ser Met Cys Lys Pro Ile Thr Gly Thr Ile Asn Asp Leu Asn Gln
 20 25 30
 15 Gln Val Trp Thr Leu Gln Gly Gln Asn Leu Val Ala Val Pro Arg Ser
 35 40 45
 Asp Ser Val Thr Pro Val Thr Val Ala Val Ile Thr Cys Lys Tyr Pro
 50 55 60
 20 Glu Ala Leu Glu Gln Gly Arg Gly Asp Pro Ile Tyr Leu Gly Ile Gln
 65 70 75 80
 Asn Pro Glu Met Cys Leu Tyr Cys Glu Lys Val Gly Glu Gln Pro Thr
 85 90 95
 25 Leu Gln Leu Lys Glu Gln Lys Ile Met Asp Leu Tyr Gly Gln Pro Glu
 100 105 110
 Pro Val Lys Pro Phe Leu Phe Tyr Arg Ala Lys Thr Gly Arg Thr Ser
 115 120 125
 30 Thr Leu Glu Ser Val Ala Phe Pro Asp Trp Phe Ile Ala Ser Ser Lys
 130 135 140
 35 Arg Asp Gln Pro Ile Ile Leu Thr Ser Glu Leu Gly Lys Ser Tyr Asn
 145 150 155 160
 Thr Ala Phe Glu Leu Asn Ile Asn Asp
 165
 40